

CEER-E-008

CEER-8

September 1978

OBSERVATIONS ON SCHISTOSOME INFECTIONS

(OF BIOMPHALARIA GLABRATA AND AN

INVADING POPULATION OF TAREBIA GRANIFERA

IN A SMALL STREAM IN PUERTO RICO

JM. Butler

FF. Ferguson

J. Palmer

W. R. Jobin

CENTER FOR ENERGY AND ENVIRONMENT RESEARCH

UNIVERSITY OF PUERTO RICO ~ U.S. DEPARTMENT OF ENERGY

---Page Break---

OBSERVATIONS ON SCHISTOSOME INFECTIONS OF
BIOMPHALARIA GLABRATA AND AN INVADING
POPULATION OF TAREBIA GRANIFERA
IN A SMALL STREAM IN PUERTO RICO?

By

R. Palmer?,

S. M. Butler?, F. F. Ferguson?,

and W. R. Jobin?

eS

*contribution of Human Ecology Division, Center for Energy and
Environment Research, Caparra Heights Station, Puerto Rico,
00935.

*Former Parasitologist San Juan Laboratories, Center for
Disease Control, US PHS.

Former Chief, San Juan Laboratories, Center for Disease Control,
USPHS.

Formerly of Division of Preventive Medicine, Health Department,
Santurce, Puerto Rico.

PHHead, Division of Human Ecology.

---Page Break---

Abstract

The purpose of this study was to monitor the presence
of cercariae of *Schistosoma mansoni* in 9 small stream in

Puerto Rico and to determine the distribution of the inter-
mediate snail host, *Biomphalaria glabrata* as well as a thiarid
snail, *Tarebia granifera*. The second snail expanded from its
original limited distribution soon after the project started

in February 1964. Within five years, the *T. granifera* had
occupied the central portion of the stream and the *Biomphalaria*
glabrata population remained only in the upper and lower reaches
which were not occupied by *T. granifera*, suggesting some form
of competition.

These observations suggest that the thiarid snail may have some value in a schistosomiasis control program. However the risk of transmitting the oriental lung fluke must be evaluated first since *I. granifera* does transmit this parasite in the Orient.

The occurrence of cercariae at two stations along the stream showed no seasonal pattern but appeared to follow a random distribution.

---Page Break---

Jarebia granifera, a small thiarid snail recently introduced to Puerto Rico from Hawaii, now occurs in almost every fresh-water body in Puerto Rico, often in enormous numbers

---Page Break---

OBSERVATIONS ON SCHISTOSOME INFECTIONS
OF BIOMPHALARIA GLABRATA AND AN INVADING POPULATION
OF TAREBIA GRANIFERA IN A SMALL STREAM IN PUERTO RICO

by

3. M, Butler,

Ferguson, 3. R. Palmer

and W. R. Jobin

INTRODUCTION

It has been observed for some time in Puerto Rico that the thiarid snail, *Tarebia granifera* was spreading throughout the island, especially in flowing water. Furthermore it often appeared to displace existing populations of *Bionphalaria glabrata*, the intermediate snail host of schistosomiasis.

During a field study on fluctuations in the numbers of schistosome cercaria produced by a colony of *B. glabrata* in a small stream,

it was noted that a limited population of *T. granifera* was

gradually expanding into the area occupied by the *S. glabrata* colony. It was the purpose of this study to record the ensuing distribution of these two snail species, as well as the fluctuations in the number of cercariae proceeding from the *S. glabrata*.

The study was conducted on a small stream on the northeastern

coast of Puerto Rico. The stream, called Quebrada Miles, is a tributary of the Quebrada Angela in the watershed of the Rio

Herrer;

It drains a small part of a community called Malpica

in the Cienega Baja area of Rio Grande, Puerto Rico (Figure 1).

---Page Break---

FIGURE LOCATION OF QUEBRADA MILES IN RIO GRANDE, PUERTO RICO

TO SAN JUAN ?10 FAJARDO

---Page Break---

MATERIALS AND METHODS

After a reconnaissance survey in February 1964, permanent sampling stations were established along Quebrada Miles, beginning with station 1 at the upstream origin and continuing downstream at 100 foot intervals for 10,000 feet. These stations were inspected about twice a year for snails, making 10 sweeps with a wire screen dipper at each station.

Because of the continuous presence of human feces and

glabrata at Stations 16 and 23 they were monitored two days

@ week for over two years to measure the number of schistosome cercariae in the water. At Station 16 a centrifuge was used to concentrate cercariae from 20 liter samples of stream water beginning in February 1965 (Butler et al, 1971). The samples were taken hourly from 10 AM to 2 PM on Tuesdays and Thursdays. The cercariae recovered were fixed with picric acid and formalin and examined with a low power microscope to determine the species. At Station 23 a filter of 11 centimeter diameter was used,

recovering the cercaria on \$ and S 404 filter paper (Rowan, 1965), After staining with ninhydrin and heat fixation the cercaria were examined microscopically to assure correct identification, The same sampling schedule was followed as at Station 16, beginning in September 1964. Local rainfall and stream flow records were obtained from the U.S. Weather Bureau and the U.S. geological Survey.

---Page Break---

RESULTS

The results of the cercarial sampling showed no definite seasonal trends (Table I). The number of cercaria recovered each day varied from 0 to 580 in the 20 Liter samples. Comparisons of cercarial populations with rainfall or water temperature failed to show significant trends, either with the number of cercariae recovered on a given day or with the number of days per month when cercariae were recovered. Most of the cercariae were recovered at 12 noon and 1 PM but significant numbers were also recovered at 10 AM, 11 AM and 2 PH. The results indicated sporadic transmission, probably related more to occasional defecation upstream than to changes in seasonal conditions. The centrifuge was easier to operate than the filter when the water contained a turbidity above 50 standard units since the filter process required additional filter papers

or pre-flocculation of the sample.

The results from the bi-annual snail surveys showed a marked interaction between the populations of *Tarebia granifera* and *Biomphalaria glabrata*. In addition scattered populations

of *Physa cubensis*, *Drepanotrema hoffmani*, *Lymnaea cubensis* and

Ferrissia beaus were observed but showed no relation to the other snail population except that *Physa cubensis* often

occurred at the same stations as *Biomphalaria glabrata*.

---Page Break---

v

TABLE 1

Ratios of numbers of days when cercaria of *Schistosoma mansoni* were recovered from 20 liters of sample versus number of days sampled each month at two stations along a small stream, Quebrada Miles, in Cienaga Baja, Puerto Rico.

eS

MONTH 1964 1965 1966

Sta. 23 Sta. 16 Sta. 23 Sta. 16 Sta. 23

2 1s 218 18

F a7 us 218 28

4 ws 9 2/10 4/10

A 9 oo o/s 218

" 218 os oo o/9

3 8/9 319 oo o/s

3 619 s/o ore o/s

a or 219 oo 9

s 24 219 a9 o/9 ve

° 49 1/8 5/8 o/s os

N ore 19 5/9 o/s o/s

o o/10 219 219 019 o/9

---Page Break---

During the first snail survey of June 1966, *B. glabrata* was present at 53 of the 100 stations from station 4 to 93 (Figure 2).

granifera was found from station 30 to 47 and at station 70, @ total of 17 out of 100 stations. Gradually

the *I. granifera* population extended downstream, reaching station 64 by April 1965 and station 88 by January 1968. Finally at the end of the study in October 1969 the thiarid Population had reached station 92.

In contrast, the colony of *8. glabrata* began decreasing in extent by April 1965, showing a marked withdrawal in March 1966 to the reach between stations 11 and 44. In 1967

glabrata was re-established in the lower reaches below station 85. During 1968 until the end of the study in 1969, *8. glabrata* appeared sporadically in the interaediata reaches.

The changes in extent of *8. glabrata* and *I. granifera* showed a strong relationship. As the thiarid snail covered the middle reaches of the stream, the planorbid snail population remained only in the upper and lower reaches, retreating from its original widespread distribution. This was seen most dramatically during October 1967 when *8. glabrata* did not occur at any of the stations from 29 to 85 where *T. granifera* was present (Figure 2).

---Page Break---

JUN? 64 OCT" 64 JAN'6S APR'®5 MAR'6S FEB'67 OCT'6? JAN'68 JAN'69 OCT? 69

CO

(em Torebiagnifera

Figure 2 Presence of Biompholariaglabrats and Torebiagrenifera along & small stream,
?Quebrada Miles, in Ciénaga Baja, Puerto Rico,

---Page Break---

This exclusion was not complete during other surveys but the relationship was marked during the last 4 years of the study. Due to the sporadic transport of snails by flow and their continual up-stream migration against the current, the Populations were developing in a dynamic situation. Thus both species are being continuously reintroduced by this process throughout the stream's length. Because of this, long-term absence of the planorbid from the middle reaches must have been due to a significant force existing in those reaches which Prevented normal development of the planorbid colony, probably the presence of *I. granifera*. A second more complicated expla-

Qranifera does not have

exactly the same habitat preferences as *B. glabrata*, changes in

nation is also possible. Since *I.*

the ecology of the middle reaches could have favored the thiarid snail and suppressed the other. Although no obvious physical changes occurred during the study, there may have been changes in flow of the stream, flood frequency and other factors which

were not monitored,

---Page Break---

10

DIScuss1ON

Over 3,500 *I. granifera* from eastern Puerto Rico were examined for trematode infections and found negative (Lee and Berrios, 1973). However, *Tarebia granifera* is the intermediate host for *Paragonimus* and theoretically could cause human disease in Puerto Rico, if un-cooked crustacea were eaten locally. This does not occur in Puerto Rico but its possibility should be

evaluated. If it were proven that *Tarebia granifera* could be used to prevent schistosomiasis however, the extremely small risk that it might some day transmit *Paragonimus* would seem acceptable. Furthermore the snail is present in many parts of the island already and its introduction into schistosomiasis control would not dramatically change the remote possibility of *Paragonimus* transmission in the future.

Carefully designed field trials are needed to determine if *T. granifera* can regularly control *S. glabrata* and to determine the habitat characteristics in which this method would succeed. Assuming that *Tarebia granifera* would be most effective in flowing water, it would complement the use of *Marisa cornuarietis* in standing water (Jobin, et al, 1973).

The data on cercarial populations did not show any particular seasonal distribution. Previous reports on monthly variations in infection rates of snails have also shown a fairly uniform seasonal pattern. For four years the Puerto Rico Health Department collected snails from their pilot control projects and checked them for a week for shedding of cercariae, in the U.S. Public

---Page Break---

an

Health Service Laboratory. The Annual Reports of the USPHS
Laboratory showed a mean infection rate of 1.5% for the snails,
varying between 0% and 5.6% but no consistent annual pattern

was found (Table 11).

---Page Break---

*parsaqtoo styeus 40 soqunu/'

ous pan994UT 40 s99urte

sez | ove fore | vez | so [sot | eco [zoo | eet] coe | ort | or

mo:

esneyee | vate/es | eroeres | eeceree | vascise | nesecce | lezcerer | soce/z9 | zozovert | Joceo/czt |
rersies | fesea/zot

o | two [sez | oo | eee | tor

ose

seere | fercryan | reste | eserce | loserer

"0 wr [et [ows [ee [seo | cco | ovo | oe | use | wo | ot

966

Seurse | tserer | eserst | soorson | seers» fuerte | Jocotr9 | sosziz | josseree | Joserris | custior | secret

260 wet {cco | foweo | cer [et | to | eet | oo | ost | 900 | ore

ss

zoaistt | susyor} | ooever | ssztre | zeersee | evrrver | ceers | sesevce | leerse | foscarrer | esoz/er
jocse/ne

mez [owme [ewe [acct [acco | faut | ero [wo | wo wee | wero [asro

86

czeryen | s9/se | scores | ozorsot | evorss | enerst | sos | w92/0 | escio | ezevoe | onsie | ensoyt

faaqeoooq | aaquanoy | soqoaq | -ada5 | asnby | Aire | auc} fea} | tracy | voaew [Axenaqag |

Anemueg] aeos

"966 BNC HONOWL SST AUVNNE Kou - VIVID WIV WAGIOTS 40

SNOTL237I00 WoO MOUS NSMVL VINO ALTATIOBANT ~ TIWNS NWOTY O14Ne 40

AUWHANS:

1 eve

---Page Break---

rr

Acknowledgnents

Acknowledgment Is made of the assistance of Hr. Ramin Méndez and Marta Luisa Santos of the Puerto Rico Health Departaent and Ne. Antbal Carrign of the US Public Health Service for performing

the majority of the field work.

---Page Break---

*

*

1?

References:

A, Butler, G. M., Rutz-Tiben, E., and Ferguson, F. F. 1971

?Evaluation of two methods for the detection

of Schistosoma

Sansonii cereariae shed by Biomphalaria glabrata". A.3.1.M.M.

V 20, 157-159.

2. Rowan, B. 1965 "The ecology of schistosome transmission

foci" Bull WHO, V 33, 63-71.

3+ Lees O, and Becrios-Duran, L. 1973, "Influence of Freshwater

Trenatodes in Eastern Puerto Rico on Schisto:

asis Skin

Test Results" Health Service Reports, V 68, 878.

0.

4 Jobin, W. R., Ferguson, F. F. and Berrios-duran, Le A. 1973

?Effect of Marisa cornuarietis on populations of Biomphalaria
alabrata in farm ponds of Puerto Rico" V 22, 278-266.

---Page Break---