

support.

Corrected Text:

Paper A127 - 'TEMPORAL PATTERNS OF THE METAZOAN PARASITES IN THE WHITE MULLET, (MUGIL CUREMA VALENCIENNES) FROM JOYUDA LAGOON, PUERTO RICO' by Jorge R. Garcia and Environmental Center for Energy and Environment Research

TEMPORAL PATTERNS OF THE METAZOAN PARASITES IN THE WHITE MULLET, MUGIL CUREMA VALENCIENNES, FROM JOYUDA LAGOON, PUERTO RICO* by Jorge R. Garcia. This work was partially supported by The Marine Ecology Division of the Center for Energy and Environment Research, University of Puerto Rico, Mayaguez. MS Thesis, Marine Sciences, RUM UPR(1981)

Abstract: Monthly prevalence, incidence, and intensity of metazoan parasites from the white mullet, *Mugil curema Valenciennes*, were examined from February 1979 to April 1980. Variations in the prevalence and incidence percentages displayed by external parasites suggest patterns of periodical occurrence. The periodicity of different parasitic populations is related to the changes in relative abundance and sexual maturity of mullet, short-term salinity variations, and interspecific associations of parasites. Four new host-parasite associations were noted in *Mugil curema*: *Pseudohaliotrema mugilinus* Hargis, 1955, *Metamicrocotylea macracantha* Alexander, 1954, *Lernaeenicus longiventris* Wilson, 1917, and *Trachelobdella lugubris* Grube, 1840.

Acknowledgements: I am indebted to the Marine Ecology Division of the Center for Energy and Environment Research for providing the cooperative personnel and research facilities for this investigation. I am especially grateful to Dr. Ernest H. Williams, Dr. José M. López, Dr. Luis R. Almodovar, and Dr. Paul Yoshioka for the revision of this work. I would also like to acknowledge the technical assistance provided by Carlos Bonafé, Edwin Gonzalez, Pablo Cabassa, and Miguel Nieves. Also for Roberto Castro, Mario Perez, and Luis Negron for their cooperation and interest in this investigation. Finally, but most importantly, I am very grateful to my parents and to my wife for their economic and moral support.

Support.

ACKNOWLEDGMENTS.

LIST OF FIGURES.

LIST OF TABLES.

INTRODUCTION.

LITERATURE REVIEW.

MATERIALS AND METHODS.

The Study Area.

Sampling Procedures.

Hydrological Measurements.

TABLE OF Laboratory Procedures.

Statistical Analysis.

RESULTS AND DISCUSSION.

Courses Hydrological Parameters.

Relative Abundance of White Mullet.

Host - Parasite Interactions in Joyuda.

Interactions Between Parasitic Species.

Parasite-Ecosystem Interactions.

CONCLUSIONS.

BIBLIOGRAPHY.

PAGE

2

2

2

15

1

2

28

6

Figure List:

Figure 1: Joyuda Lagoon and adjacent coastal features. Page 10

Figure 2: Monthly means and range of salinity measurements at Joyuda Lagoon, between the period of February 1979 through April 1980. Page 6

Figure 3: Monthly means and range of water temperature measurements at Joyuda Lagoon, between the period of February 1979 through April 1980. Page 2

Figure 4: Mean and range of dissolved oxygen values at Joyuda Lagoon for the period between February 1979 and April 1980. Page 8

Figure 5: Relative abundance of white mullet in monthly collections at Joyuda Lagoon with 95% C.L. (number in parenthesis represents sample size). Page 21

Figure 6: Standard length distribution of white mullet in monthly collections at Joyuda Lagoon, for the period between February 1979 and April 1980. Page 24

Figure 7: Monthly variation in the incidence percentages of the Acanthocephalian parasite, *Florida mugilis*, occurring in the white mullet. Page 27

Figure 8: Monthly variation in the incidence percentages of the monogenean trematode *Pseudoshaliootrema mugilis*, parasitic in the white mullet. Page 27

Figure 9: Monthly variation in the incidence percentages of the monogenetic trematode, *Metanicotyle macracantha*, parasitic in the white mullet. Page 28

Figure 10: Monthly variation in the incidence percentages of the copepod, *Ergasilus lizae*, parasitic in the white mullet. Page 28

Lernacenicus longiventris, a parasite on the white mullet. This is followed by a discussion on the monthly variation in the incidence percentages of the copepod, *Bonolochus coninmus*, parasitic in the white mullet.

Figure KITT illustrates the monthly variation in the intensity of parasitization of the copepod, *Ergasilus Liza*, on the white mullet. The following figures, Figure XIV and Figure XV, show the monthly variation in the intensity of parasitization of the monogenetic trematode, *Metenicocotylea macracantha*, and the copepod, *Lernacenicus Longiventris*, on the white mullet respectively.

Table 1 to Table IX provide various data. Table 1 lists the parasites of the white mullet, *Mugil curema*, in Joyuda Lagoon. Table II presents the monthly distribution of salinity (‰), temperature and dissolved oxygen (p.p.m.) values from Joyuda Lagoon during the period between February 1979 and April 1980. Table III presents the index of gonad development in *Mugil curema* for the same period. Table IV shows the monthly prevalence of parasitic species in the white mullet from Joyuda Lagoon during the same period.

Table V is a contingency table representing the negative interspecific association between the monogenetic trematode *Metamicrocotylea macracantha* and the copepod *Ergasilus liza*. Table VI represents the positive interspecific association between the monogenetic trematode *Pseudochaliootrema mugilinus* and the copepod *Bonolochus coninmus*.

Table VII and Table VIII analyse simple linear regressions between monthly means of salinity and temperature, respectively, and intensity of parasitization by metazoan species of parasites in white mullet.

Table IX presents a multiple linear regression analysis between monthly means of salinity and temperature and intensity of parasitization by metazoan species of parasites in white mullet.

Parasitization by metazoan species of particles in white mullet...

[INTRODUCTION] The importance of ecological studies in fish parasite populations has been increased by the development of aquaculture programs with freshwater and marine fishes. Artificial culture conditions enhance the direct transfer of some parasitic species and usually reduce the natural defense mechanisms of the hosts against parasitic infections. Knowledge about fish-parasite interactions in their natural ecosystem provides useful baseline information for aquaculture research. The present study was conducted to provide such baseline information in Fish which has potential for aquaculture. The specific objectives in this study were:

- 1) To study the periodic occurrence and intensity of infection by metazoan parasites in the white mullet, occurring in Joyada Lagoon.
- 2) To examine the influence that changes in the external environment have on the parasitic fauna of the white mullet.
- 3) To determine the effect that spawning and possible migrations of the fish have on its parasitic composition.
- 4) To add some ecological and biological information about the white mullet.
- 5) To present a general description of the study site, Joyada Lagoon, and to evaluate the role of this particular ecosystem in the life cycle of the white mullet.

The White mullet is the most abundant among four species of mullet reported in Puerto Rico. A typical inhabitant of estuaries and brackish-water systems, the white mullet is the most common representative of gray mullet in the Caribbean coast. Anderson (1957) described the early development, spawning, growth and occurrence of the white mullet along the south Atlantic Coast of the United States. Yanez-Arancibia (1976) studied the feeding habits, growth, spawning behavior and trophic relations of the white mullet in the lagoonal system of Guerrero, on the Pacific Coast of Mexico. Because of its size and abundance, the white mullet, locally known as "jarea", is a prominent commercial

Fish in Puerto Rico. Besides being a common food item in some parts of the island, mullet are highly prized as bait for sport fishing. White mullet are especially important because of their potential for culture (Yashouy 1972, Tang 1975, Sebastian and Nair 1975, De Silva and Wijeyaratne asm).

LITERATURE REVIEW

The general biology and ecology, as well as some parasitological information, is available concerning the striped mullet, *Mugil cephalus*. This species has gained much attention since it is

more broadly distributed and commercially important in the United States (Anderson 1957). Rawson (1976) studied the population biology of parasites of *Mugil cephalus* in a temperate area and found that the number of parasite species increased with the age of the host and that initial infection was influenced by the close association of mullet age-classes. Skinner (1975) surveyed the parasites of *Mugil cephalus* in Biscayne Bay.

Recent studies have approached some ecological aspects of fish parasite populations in natural ecosystems. However, most of this work has been done in subtropical and temperate areas. Overstreet (1968) calculated significant correlations between monthly means of temperature, salinity, and size of the fish with the mean number of parasites infecting the inshore Lizardfish, *Synodus foetens*, from an estuarine canal in South Florida. Williams (1979) studied the seasonal incidence of the cestode *Isoglaridacris viscogensis* in its fish host, *Ilypentelium nigricans*, and associated low incidence of the parasite with cold winter temperatures and decreased feeding by the fish. Evans (1978) described an annual cycle of occurrence and maturation in the "roach" *Rutilus rutilus* by the cestode *Asymphyldora kubanicum* in the Worcester-Birmingham channel in England. Meskal (1966) reported seasonal variations of parasites from the cod in coastal waters of

Norway and related fluctuations in the trematode population with the seasonal abundance of a copepod, apparently acting as an.

Intermediate host. Grimmes and Miller (1976) suggested a "temperature dependent rejection response" as a possible factor involved in the seasonal periodicity of three species of caryophyllaeid cestodes in the creek chubsucker, *Erimyzon oblongus*, in North Carolina. Rawson (1972) reported seasonal abundance of monogenetic trematodes in the bluegill, *Lepomis macrochirus*, in a reservoir in Alabama. Other studies demonstrate that differences in the habitat and geographic location are more important than seasonal variables in the parasitic composition of some fishes (Schroeder 1970, Dowgiallo 1979). Paperna (1975) stated that mortalities of gray mullet cultured in brackish and freshwater ponds, which have resulted from massive infections by parasites, appear to be coupled with environmental stresses such as declines in oxygen, critical changes in temperature or salinity, or pollution. Taxonomical references on parasites infecting the white mullet are scattered in the literature. Table 1 presents a list of parasites which have been noted previously.

Materials and Methods: The study area, Joyuda Lagoon (Figure 1), is a coastal brackish water system located on the southwestern coast of Puerto Rico. The lagoon is approximately 1.6 km long and 0.8 km wide, covering an area of 121 hectares. A conspicuous band of the red mangrove, *Rhizophora mangle*, borders the lagoon. There is also some development of the black mangrove, *Avicennia nitida*, and white mangrove, *Laguncularia racemosa*, on the western section. Water exchange between the lagoon and the sea occurs along a small channel bordered by mangrove which opens seaward into a sandy patch area of turtle grass with scattered coral growth. The average water depth of the lagoon is 1.3m, with a maximum depth of approximately 4m. The bottom substrate is composed of soft mud sediments.

The text contains organic detritus, primarily derived from mangroves. **Sampling Procedures:** Monthly samples of ten mullet from Joyuda Lagoon were examined for metazoan parasites from February 1979 through March 1980. Most of the collections were made with monofilament and

nylon bottom gill nets 60 m long and 2.5 m high, with a square mesh diameter of 3.9 cm. Some individuals were captured with cast nets. All the sampling was done from a 136" fiberglass boat powered by a small outboard motor. The nets were set at sunset and recovered at dawn. Fish were placed in individual plastic bags and taken directly to the laboratory for examination.

Figure 1, Joyuda Lagoon and adjacent coastal features 10

Suana Sub River Bore Som El z= 620m Rincon = Lagoon Ta. Coastline Lagoons Ed Mangrove
Scale 1:100,000 NOAA Nautical Chart 901

Salinity, temperature, and dissolved oxygen measurements were obtained every month at five stations in the lagoon. These data were recorded on sampling days at the time of setting the nets with YSI meters from Yellow Springs Instruments Co. Model 33 S-C-T was used to measure salinity (to the nearest + 0.5%) and temperature (to the nearest + 0.6 °C). Model 57 was used to measure dissolved oxygen (to the nearest ± 0.2 ppm). The instruments were calibrated according to the instruction manual at least once a month.

Laboratory Procedures: The fish samples were taken directly from the field to the laboratory facilities at CEER and examined within a few hours of collection. All fish were weighed and measured (standard length), and the gonads were removed from the fish and weighed. Parasitological examinations were limited to metazoan (multicellular) species and included the body surface, gills, and alimentary tract of the fish. Gill arches were removed and placed in separate dishes. Numbers and species of parasites present were recorded for each gill arch. An incision was made on the right side of each fish and the different organs were examined.

In the statistical analysis, the incidence percentage and intensity of infection by the different parasitic species were noted as monthly values for the 15-month period. Incidence percentages represent the proportion of fishes infected by a particular parasite. Intensity of infection values represent the proportion between the total number of individuals of parasitic species and the total number of mullet examined in a month. A logarithmic transformation ($\log N$) was applied to the monthly values of intensity of infection. Prevalence represents the presence of a parasite in a monthly sample. Simple and multiple regression analyses between the temperature and salinity means and the intensity of infection values were calculated for every species except for the leech, *Trachelobdella Lubrica*, which only occurred once during the study period. The correlation coefficients were obtained from a program of an Apple I computer, which also calculated the standard error of estimate. Possible interrelations between parasitic groups were tested for.

RESULTS AND DISCUSSION

Hydrological Parameters

Salinity - The average salinity in Joyuda Lagoon for the period between February 1979 and April 1980 was 19.9 ppt. Monthly mean values ranged between a minimum of 12.0 ppt in October 1979

to a maximum of 30.0 ppt in April 1980. Figure 1 presents the monthly fluctuations in salinity during the study period. Summer months (May-October) averaged lower salinity values (avg 18.6) while the winter period (November-April) presented a higher average value (avg 20.8). Monthly mean values indicate that the salinity pattern in the lagoon is unstable and that moderate variability can occur in short time intervals of less than one month. Salinity is mostly determined by the amount of precipitation and runoff, the temperature, wind effect on evaporation, and the intrusion of seawater during high tides.

Temperature - The average water temperature was 27.7°C. Monthly mean values ranged between a maximum of 30.0°C in May and July 1979 and a minimum of 24.7°C in February 1980. Figure 2 presents the monthly variation in mean temperature values. The average water temperature was higher during the summer months.

(May-October) with an average of 20.8°C as compared to the winter months (November-April) with an average of 27.0°C. The gradual decrease of water temperature started in September and reached its lowest point by February. The pattern of water temperature is affected by air temperature because of the shallow nature of the lagoon and relatively stagnant condition of the water.

Short term variation in water temperature may occur as a result of heavy precipitation and the drain of cold freshwater from the runoff of adjacent mountains. Dissolved oxygen - the content of dissolved oxygen in the water ranged from 4.7 ppm in March 1980 to 7.2 ppm in April and December 1980. Although moderate variation occurred on a monthly basis, the summer period registered a rather stable content of oxygen in the water compared to the winter months in which the degree of variation from month to month was high (see Figure IV). If we account for the temperature and salinity effect on the different expected levels of oxygen saturation, the variation in the monthly percentages of oxygen saturation must be related to biological processes occurring in the lagoon. Table 17 presents the monthly values of the percentages of oxygen saturation.

Relative Abundance of White Mullet in Joyuda Lagoon: The relative abundance of white mullet in monthly samples indicated a peak in February 1979 with 41% of the total capture in a sample size of 193 individuals. Another high value was recorded in September 1979 with 32% in a sample size of 49 individuals. Figure V presents the monthly variation in relative abundance of white mullet. The sudden decrease in abundance of white mullet after February tends to support the.

The theory of an offshore spawning migration of this fish has been suggested by Anderson (1957), Moore (1974) and Yanez (1976). The index of gonad maturity (see Table IIT) indicated that 80% of the mullet examined in February 1979 had an advanced stage of gonad development. In March 1979, all the individuals examined were sexually mature. The following months presented some mature individuals.

Unintelligible text.

Unintelligible text.

Unintelligible text.

Unintelligible text.

As in the collections until September, no individuals were found to be sexually mature. The peak of adult white mullet in September with reduced gonad development may be indicative of a return, in part, of the white mullet population after the spawning had taken place in offshore waters. The overlapping date for the months of February and March 1980 are not significant in this matter because the individuals examined were not adult fishes (see Figure VI). The low values in relative abundance after October 1979 may be indicative of the detrimental effect of hurricanes David and Frederick in September 1979, or due to over-fishing of the adult mullet population in the lagoon during their spawning season.

Period of prevalent spawning in February and March 1979 by local and commercial fishermen. Host = Parasite Interactions. Six species of metazoan parasites were found to infect the white mullet during the study period at Joyuda Lagoon. Temporal patterns of incidence percentages must be interpreted with caution due to small sample size numbers. The Acanthocephalian parasite, *Floridosentis usitis* (Machado Filho 1951), which is an internal parasite, localized always in the intestine of the fish, was present at least once in every monthly sample. *Floridosentis mugilis* registered a peak of incidence in monthly collections of March 1979, and then in December 1979, January and March 1980. Figure VII presents the monthly fluctuations in incidence percentages for this species. The pattern of incidence percentages does not seem to be directly determined by any external factor related to the water quality of the lagoon. The fact that *F. mugilis* is transmitted by an intermediary host is indicative.

26 of the apparent availability of this intermediate host throughout most of the year in the lagoon. Further studies on the life cycle can inform about the short term variability in the incidence percentages of this species. Two monogenean trematodes parasitized the gill filament of the white mullet in Joyuda Lagoon. *Pseudohaliotroma mugilinus* Hargis 1955 presented peaks in incidence percentages in February and March 1979, and then in March and April 1980. This monogenean parasite was absent from August to October 1979. Nevertheless, it appeared with moderately high incidence percentages throughout the study, especially during the winter period (Figure VIII). Low salinities, or perhaps the sudden decrease in salinity from August to September, continuing into October 1979, could have caused the absence of this parasite during these months.

Metanicrocotyle macracantha Alexander 1954

Another monogenetic trematode, which infects the gill filaments, peaked in November 1979 and March 1980, and was also common in July and September 1979 and February 1980. *Metanicrocotylea gacracantha* can withstand short time salinity variations and was present in September 1979 (see Figure 2x) when salinity reached its lowest point during the study (approx. 12 ppt). The presence of this monogenetic trematode in *Mugil curena* was first noted in May 1979, but continued to appear in the rest of the study. This strongly suggests that the parasite either entered

the lagoon after a sand bar formation opened in April-May with early spawned adult females, or with juvenile mullet which were probably new recruits in the lagoon.

However, after May, the short-term variations of incidence on the host may be related to other factors, either environmental, biological, or both. Three species of copepods of different genera were found on the white mullet. *Ergasilus lizeo* Kedyer 1864, which occurred only in the gill filaments of the fish, was more abundant in February 1979. It prevailed until March 1979 and then disappeared until October 1979, reaching another high incidence value during January 1980 (see Figure X). Although *E. Ligae* occurred during the winter period, its incidence in the white mullet seems to be more related to the migratory pattern of the adult fish than to seasonally related hydrological conditions. *Ergasilus Ligae* survived in a wide range of temperature and salinity variations in Joyuda Lagoon. Its absence after March 1979 could be due to the migration of adult mullet to offshore waters.

As a consequence of this migration, the individuals examined during the remaining summer months did not possess this parasite because they were new recruitments composed mostly of juvenile fish. With the next immigration of adult mullet after September 1979, the parasite became again a regular component of the parasitic population of the mullet in Joyuda Lagoon.

The parasitic copepod, *Lernacenicus longiventris* Wilson 1917, was found partially embedded in the fins and body surface of the white mullet. Its prevalence and incidence percentages show a clear peak during summer with consistently high incidence percentages for the months of July, August and September 1979 (see Figure XI). After a lag period of two months, the parasite appeared again in the collections, reaching another peak during March 1980.

The pattern of occurrence of this parasitic species is again indicative of the different populations of mullet which were sampled during the study period. The parasite first appeared in monthly collections from June 1979, one month after the migration of the fish, and then showed reduced incidence percentages in October 1979, probably as a result of uninfected adult mullet examined in that particular monthly sample.

Apparently, after being introduced by new recruitment into the lagoon during the summer period, the parasite adapted well to the strongly variable hydrological conditions in the lagoon and persisted in the samples despite a change of 17 ppt in salinity between October 1979 and April 1980.

Bomolochus concinnus Wilson 1911 occurred mostly in the mucus of the branchiostegal cavity with occasional presence in the branchiostegal filaments. *Bomolochus concinnus* appeared in nine out of 15 monthly samples, but did not show any distinct peak of abundance.

The text appears to be corrected as follows:

Based on reasonable sample size numbers, its abundance during the rainy period in summer (see Figure XIX) from August to October 1979 may be indicative of low tolerance to the sudden salinity decrease associated with hurricanes David and Frederick.

Interactions Between Parasitic Species:

Six species of external parasites and one endoparasite were present in white mullet at different time periods throughout the study. The prevalence of these species in the white mullet is presented in Table IV. It was observed that ectoparasites which occupied similar microhabitats within the fish, such as *Metapicrocotylea*.

Sonar 97

Table IV: Monthly prevalence of parasitic species in the white mullet from Joyuda Lagoon, during the period between February 1979 and April 1980.

Species: Sp1, Sp2, Sp3, Sp4, Sp5, Sp6

Feb: +, -, +, +, -, .

Mar: +, -, ., +, -, +

Apr: ., -, -, -, -, +

May: +, +, -, -, +, .

Jun: +, +, ., +, ., .

Oct: -, -, -, ., +, +

Nov: +, +, +, -, -, +

Dec: -, -, -, +, -, +

Jan: +, -, +, +, ., +

Feb: +, +, ., -, +, +

Mar: +, +, -, +, -,

Apr: -, -, -, +, +, +

(- = Absence, + = Presence)

Sp. 1 ~ *Pseudohalioirens mugilinus*

Sp. 2 ~ *Notamicrocotylea macra*

Sp. 3 ~ *Bonolochus concinnus*

Sp. 4 ~ *Ergasilus lizae*

Sp. 5

Sp. 6 ~ *Floridosentis mugilis*

As *Nacracantha* and *Ergasilus lizae* did not occur together in monthly samples. Both species are specialized parasites of the gill filaments. Table V details the presence-absence display of both species in monthly samples. These species show a significant negative association ($p=0.017$). Possibly one of these parasites is excluding the other in competition for food and/or space. Their mutual exclusion in monthly samples could be affected by the seasonal periodicity of one species or the other, their adaptations to withstand environmental stress, chance, or actual interspecific interactions between both species. The association between *Pseudohalioirens mugilinus* and *Bonolochus concinnus* also suggests possible interactions between different parasitic species.

Populations. In this case, a positive association was found between both parasites (see Table V), and the probability of association was highly significant ($p=0.002$). *Pseudochalictrena mugilinus* and *B. concinnus* are exposed to similar environmental conditions as both are parasitic in the branchiostegal cavity. However, each species occupies different niches or microhabitats within the fish. *Bonolochus concinnus* was always found in the mucus layer of the branchiostegal flap, while *Pseudohalioirens mugilinus* was always parasitic in the gill filaments. Different food and space requirements can allow both species to coexist together in the host. *Floridosentis mugilis*, an acanthocephalan worm, was the only internal parasite observed and it prevailed consistently in every monthly sample (see Table IV). The presence of other endoparasites in the alimentary tract is probably limited by the lack of intermediate

TABLE V: Contingency table representing the negative interspecific association between the monogenetic trematode *Metamicrocotyles macracantha* and the copepod *Ergasilus lizae*.
Notamicrocotylean Absent, 6 3 9 *Ergasilus Lizae*. Probability of association by chance (p) = 0.017

TABLE VI: Contingency table representing the positive interspecific association between the monogenetic trematode *Pseudochalictrema mugilinus* and the copepod *Bonolochus concinnus*.
Present absent Present ° ° 2 *Bonolochus concinnus*. Probability of association by chance (p) = 0.002

Hosts such as mollusks, which are rare in Joyuda Lagoon. The possibility of competitive exclusion by the acanthocephalan is contradicted by the presence of 2 genetic trematodes which occur along with *E. Mugilis* in collections of mullet examined in La Parguera, Puerto Rico (Williams, PH, unpublished data). The consistent patterns of association between parasites constitutes evidence that their temporal occurrence is real and not an artifact of sampling variability. Parasite-Ecosystem Interactions: Simple and multiple regression.

Analyses between monthly means of temperature and salinity and the intensity of infection by parasitic species were non-significant at the 0.05 level. Evidently, most ecto-parasitic species in *Mugil curema* withstand some degree of variation in salinity and temperature (see Tables VII, VIII, and IX). Consequently, the spread of points on both sides of the regression lines results in a high standard error and low correlation coefficients. Deterministic effects of hydrological parameters on the parasite composition of *Mugil curema* were observed for abrupt salinity variations in August through October 1979. Most species of external parasites presented low incidence or were absent during this period. The effect of short term variation is probably more important in determining the prevalence of some parasitic species than seasonally related variations which are gradually experienced by external parasites. If we consider the wide range of tolerance that mullet have for salinity variations, it is suggestive that the external parasitic fauna has also adapted to variable salinities.

The latter part of the text seems to be garbled or in a different language and is unreadable. Please provide a clear version of the text for further assistance.

Two key studies today discuss about the unique complexities surrounding system automation. It's an area of study and research that is always changing.

Firstly, it's important to understand the complexities and nuances of system automation. It's not a straightforward process, but requires in-depth knowledge and understanding.

The periodic occurrence of parasitic species such as the copepods *Lernaeenicus longiventris* and *Ergasilus lizse* is explained by the migratory behavior and relative abundance of adult mullet in Joyuda Lagoon rather than seasonally determined by temperature and/or salinity. The influence of monthly variations in water temperature and salinity may play an important role in particular interspecific interactions between two species in competition for similar habitats in the fish. For example, *E. lizse* presented higher intensity of parasitization during the winter period (see Figure XIII), while *M. macracantha* had high intensity of infection in two monthly collections in summer and also prevailed in samples from September 1979 (see Figure XIV). This suggests that *M. macracantha* is more tolerant to high temperatures than *E. Lizse* in Joyuda Lagoon. *L. longiventris* also displayed a trend of occurrence during the summer months with high intensity of parasitization in higher water temperatures than the average for the ecosystem (see Figure XV).

Continuing the examination of this topic, it's essential to delve deeper into the specifics and understand the various elements at play.

One key aspect to consider is the impact of automation on different sectors. It's not a simple process, but involves a multitude of factors and considerations.

Moving forward, it's crucial to understand the implications of these findings and how they can influence future studies and developments in this field.

In conclusion, the range of tolerance observed by parasitic species to temperature and salinity variations in Joyuda Lagoon is indicative of the adaptations that these parasites have developed in order to withstand similar environmental conditions.

The host is adapted to gradients. The migratory behavior of adult mullet to offshore waters, and their eventual return to the lagoon, accounts for much of the variability observed in the periodic occurrence of some parasitic species in white mullet. Seasonally related variations in water temperature and salinity are not related to the incidence and intensity of infection by parasitic species. Nevertheless, sudden salinity variations are detrimental to some ectoparasitic populations, which are apparently less tolerant than their host to rapid, low salinity stress. Negative interspecific interactions between parasitic species occurred between populations that occupy similar microhabitats within the host. This type of association was significant for the copepod *Ergasilus liz* and the monogenetic trematode *Metanicrocotylea rantha*. Their actual competition and mutual exclusion may explain the short-term variations that these species present in their patterns of occurrence. Joyuda Lagoon is a detritus-based ecosystem that provides high food availability and protection for juvenile and adult mullet. The population proportion of adult mullet is higher during the winter period prior to their peak of sexual maturity. This fact suggests that the mullet concentrate in the lagoon in order to feed extensively and store enough energy for their spawning migration. Joyuda Lagoon may also function as a shelter for juvenile and adult mullet during periods of high wave energy and low food availability on the coast.

BIBLIOGRAPHY

Avezcua-Linares, F. 1977. Generalidades ictiológicas del sistema lagunar costero de Huizache-Caimanero, Sinaloa, Mexico. *Anales Centro Ciencias del Mar y Limnología*. Universidad Nacional Autónoma, Mexico. 4 (1): 1-26.

Amin, O.M. 1975. Host and seasonal associations of *Acanthocephalus Parksides* Amin, 1974 (*Acanthocephala*: *Echinorhynchidae*) in Wisconsin fishes. *Journal of Parasitology* 61 (2): 318-329.

Anderson, W.M. 1957. Early development, spawning, growth and

"Occurrence of the silver mullet (*Mugil curema*) along the South Atlantic coast of the United States.

Fishery Bulletin 57 (119): 397-414, Austin, H.M, 1971. A survey of the Ichthyofauna of the mangroves of Western Puerto Rico during December 1967 - August 1968. *Caribbean Journal of Science* 11 (1-2); 27-39. and S. Austin. 1971. The feeding habits of some juvenile grazing fishes from the mangroves in Western Puerto Rico, *Caribbean Journal of Science* 11 (3-4): 171-171.

Amolé, E. and J.R. Thompson. 1958. Offshore spawning of the striped mullet, *Mugil curema*, in the Gulf of Mexico. *Ichthyological notes, Copeia* (2); 130-132. Beneden, P.J. 1870. The fishes of the coasts of Belgium, their parasites and their commensals. *Bulletin de l'Academie Royale de Sciences de Belgique* 1 (1): 51-64. De Silva, B.S. and Y.J.S. Wijeyaratne. 1977. Studies on the

biology of young gray mullet, *Mugil cephalus* (L) 12. Food and feeding, *Aquaculture* 12: 159-167.

Dossiel, V.A., C.K. Petruchevski and Y. Folyanski. 1958. *Parasitology of Fishes*. Leningrad University Press, 1958. English translation Oliver and Boyd. Edinburgh and London, 1961.

Dowgiallo, M.J. 1979. Variation of metazoan parasites of the French grunt, *Haemulon flavolineatum* (Desmarest) (Osteichthyes: Pomadasyidae), by habitat type and season with an analysis of competition among parasites. Master Thesis, Department of Marine Sciences, RUM, Mayaguez, Puerto Rico.

Evans, N.A. 1977. The occurrence of *Sphaerostoma bramae* (Digenea: Allocreadiidae) in the roach from Worcester – Birmingham canal, *Journal of Helminthology*. 51: 189-196.

Evans, N.A. 1977. The site preferences of two digeneans, *Asymphylogora kubanicum* and *Sphaerostoma bramae*, in the intestine of the roach, *Journal of Helminthology*. 51: 197-204.

1978. The occurrence and life history of *Asymphylogora kubanicum* (Platyhelminthes: Digenea: Monorchidae) in the Worcester-Birmingham canal, with special reference to the feeding habits of the definite host, *Rutilus rutilus*, *Journal of Zoology*, London. 194; 143-153.

Grimes, L.R. and"

Miller, G.C. 1976. Seasonal periodicity of three species of *Coryophyllacia* cestodes in the creek chubsucker *Erimyzon oblongus* (Mitchill) in North Carolina. *Journal of Parasitology* 62 (3): 434-441.

Hopkins, S.H. 1954. The American species of trematodes confused with *Bucephalus* (*Bucephalopsis*) *haineanus*. *Parasitology*. 33 (3-4): 353-370.

Meyer, W. 1863. Bidrag til kundskab om snyltekrebsenes naturhistoriskes tidsskrift. 2(1-2); 75-320.

Machado-Filho, D.A. 1951. Una nova especie do genero *Atactorhynchus*, Van-Cleave, 1935. (*Acanthocephala*, *Neoechinorhynchidae*). *Revista Brasileira de Biologia* 11: 29-31.

McErlean, A.J., O'Connor, S.G., Mihursky, J.A. and Gibson, C. 1973. Abundance, diversity and seasonal patterns of estuarine fish populations. *Estuarine and Coastal Marine Science* 1: 19-36.

Meskal, P.H. 1966. Seasonal fluctuations in the population of two common trematode species from the stomach of the cod. *Sarcia* 26: 13-20.

Moore, D., Brusher, H-A., and Trent, I. 1970. Relative abundance, seasonal distribution and species composition of demersal fishes off Louisiana and Texas, 1962-1964. *Contributions in Marine Science* 15: 45-70.

Moore, R.I. 1974. General ecology, distribution and relative abundance of *Mugil curema* and *Mugil cephalus* on the South Texas coast. *Contributions in Marine Science*. 18: 241-255.

Manter, H.W. and Cable, R.M. 1963. Digenetic and *Aspidogastrea* trematodes from marine fishes of Curacao and Jamaica. *Tulane Studies in Zoology*. 11 (4): 169-224.

Orange, C.J. 1961. Spawning of yellowfin tuna and skipjack in the Eastern Tropical Pacific, as inferred from studies of gonad development. *Inter-American Tropical Tuna Commission*. 5(6): 461-463.

- Overstreet, R.M. 1968. Parasites of the inshore lizardfish *Synodus foetens* from South Florida, including a description of a new genus of cestodes. *Bulletin of Marine Science* 18 (2): 445-469.
- Pagan, F.A. and Rustin, H.M. 1970. Report on a fish kill at Laguna Joyuda, western Puerto Rico, the summer, 1967. *Caribbean Journal of Science* 10 (3-4): 203-208.
- Faperna, I., 1975. Parasites and diseases of gray mullet (*Mugilidae*) with special reference to the seas of the Near East. *Aquaculture* 5: 65-80.
- Rawson, M.V., 1976. Population biology of parasites of striped mullet, *Mugil cephalus* L. I. *Monogenea. Journal of Fisheries Biology* 9: 185-194.
- Sebastian, M.J., and V.A. Nair., 1975. The induced spawning of the gray mullet, *Mugil macrolepis* (Aguas) Smith and the large-scale rearing of its larvae. *Aquaculture* 5: 41-52.
- Schroeder, R.E., 1970. Ecology of the intestinal trematodes of the gray snapper, *Lutjanus griseus*, near Lower Matecumbe Key, Florida, with a description of a new species. *Studies in Tropical Oceanography*, Number 10, 151-223.
- Skinner, R., 1975. Parasites of the striped mullet, *Mugil cephalus*, from Biscayne Bay, Florida, with description of a new genus and three new species of trematodes. *Bulletin of Marine Science* 25 (3): 318-345.
- Tang, Y.A., 1975. Collection, handling and distribution of gray mullet fingerlings in Taiwan. *Aquaculture* 5: 81-84.
- Tate, H.W. and R.C. Clelland., 1957. *Nonparametric and Shortcut Statistics*. Interstate Printers and Publishers Inc., Illinois.
- Vost, C., 1877. Recherches côtières faites à Roscoff; Crustacés parasites des poissons. *Archives de Zoologie Expérimentale et Générale* 6: 385-456.
- Warburton, K., 1978. Community structure, abundance and diversity of fish in a Mexican coastal lagoon system. *Estuarine and Coastal Marine Science* 7: 497-519.
- Ward, Hob., 1954. Parasites of marine fishes of the Miami region. *Bulletin of Marine Sciences of the Gulf and Caribbean*, 4 (3): 245-261.
- Williams, D.D., 1979. Seasonal incidence of *Glaridacris* larvae and *ostoni* in Red Cedar River, Wisconsin. *Iowa State Journal of Research* 53 (4): 311-316.
- Williams, E.M. and J.T. Gaines., 1974. Acanthocephala of fishes from marine and brackish waters of the Mobile Bay region. *Journal of Marine Sciences of Alabama*, 2 (3): 135-148.
- Yanez-Arancibia, X., 1976. Observations on *Mugil curema* Valenciennes in natural breeding areas,

Mexico.

Alimentación, crecimiento, madurez y relaciones ecológicas. Anales Centro Ciencias del Mar y Limnología. Universidad Autónoma de Mexico, 30: 93-124, y B.S. Nugent. 1977. El papel ecológico de los peces en estuarios y lagunas costeras. Anales Centro Ciencias del Mar y Limnología. Universidad Nacional Autónoma de México, 4 (1): 107-114. Yashouv, A. 1972. Eficiencia del crecimiento de la lisa en estanques de peces. Bamidgen 24 (1): 11-25.

---Página Siguiente---