

**CEER-T-089**

CEER - T-089

TROPICAL RAIN FOREST CYCLING AND TRANSPORT PROGRAM

PHASE 1 PROGRESS REPORT

TERRESTRIAL ECOLOGY DIVISION

CENTER FOR ENERGY AND ENVIRONMENT RESEARCH

UNIVERSITY OF PUERTO RICO

SAN JUAN, PUERTO RICO

JANUARY 1981

CENTER FOR ENERGY AND ENVIRONMENT RESEARCH

---Page Break---

CER - 1-089

TROPICAL RAIN FOREST CYCLING AND TRANSPORT PROGRAM

PHASE 1 PROGRESS REPORT

TERRESTRIAL ECOLOGY DIVISION

CENTER FOR ENERGY AND ENVIRONMENT RESEARCH

UNIVERSITY OF PUERTO RICO

SAN JUAN, PUERTO RICO

JANUARY 1981

---Page Break---

2.0

3.0

4.0

5.0

6.0

7.0

8.0

9.0

## TABLE OF CONTENTS

Page No.

EXECUTIVE SUMMARY

INTRODUCTION...

PROGRAM OVERVIEW

PHASE I METHODS .

4.1 Sampling Design ...see0

4.2 Task Methods .. 8

4.2.1 Physical Phenomena .

## 4.2.2 Mineral Inventory and Energy Studies

### 4.2.2.2 Mineral Inventory

#### 4.2.2.2 Calorimetry ....

#### 4.2.2.2 Carbon Transport;

## 4.2.3 Vegetation Studies ..sceseeseessescreeceeees

### 2.1 Plot Characterization

#### "2132. Phenology

#### 4.2.3.1.1 Mineral Inventory

#### 4.2.3.1.2 Leaf Decomposition? <2)

#### 4.2.3.1.3 Fauna) Studies «1...

##### 4.2.3.1.3.1 Invertebrates; n

##### 2 Anohibians and Reptiies .

##### 4:3 Birds .

##### 4.2.3.1.3.4 Manmays 12

## RESULTS AND DISCUSSION

### 5.1 Physical Phenomena

### 5.2 Mineral Inventory and Energy Studies .

#### 5.2.1 Mineral Inventory ve...

#### 2.2.2 Calorimetry vases 0

#### 2.2.3 Carbon Transport ...0000

### 5.3 Vegetation Studies ...-...-

#### 1 Plot Characterization

#### Phenology

#### Uitterfat 2.2.7

#### Leaf Decomposition?. :

#### Studies sesseee

Invertebrates 12.2212)

Anphibians and Reptiles

Birds .

Maral?

PHASE 1 SCHEDULE «....,

ORGANIZATION AND MANAGEMENT

LITERATURE CITED ...

APPENDICES ...





a

10.

## LIST OF TABLES

Title

Nitrogen, Nitrate, and Nitrite Values from Cara  
del Indio Sampling Station .

Carbon Concentrations of Samples from Cara  
del Indio...

Overlaid and Plot species Contribution of Leag  
Fail (ranked 1-5) based on gm Dry Weight/m<sup>2</sup>/day

?Anphibian and Reptiles Species Observed in  
the Study Area During Phase I Studies «vs...

Anurans Observed Along Transects During 1980 Wet  
Season Surveys

Frequency of Occurrence of Food Items in *Anolis*  
*Sundlachi* and *A. evermanni* ss...

Average Number of Individuals, Percent Relative  
Abundances, and Minimum Population Density.  
Estimates of *Anolis* Lizards Observed During 1960  
Wet Season Surveys

Total Observations During July-November Transect  
Counts Corrected for Number of Counts and Time  
Spent Counting



ber of Captures and Capture Rate for Plots

ne

Mean Weights (in grams) of Seven Species Caught

in Mist Nets... : :

it

3

?

46

a

31

---Page Break---

## LIST OF FIGURES

Uitte

Generalized Ecosystem Modet

Cycling and Transport Program

Study Areas...

Vegetation Sampling Locations

within Study Plots...

Wet Weight/Dry Weight Relationships

for *Anolis gund*

*hi* and *A. evermanni*

Wet Weight/Dry Weight Relationship for

*Eleutherodactylus coqui*...

Height of Occurrence of Seven Birds

## Species in the Rain Forest

### Program Management Structure .

---Page Break---

#### 1.0. EXECUTIVE SUMMARY

This program was designed to fulfill the clear need for a comprehensive understanding of major cycling and transport processes in rain forest ecosystems so that the ecological consequences of potential energy development alternatives can be predicted. As a result of reduced OOE funding for fiscal year 1981, the original program plan (June, 1980) has been modified. The basic approach has not been altered, but some tasks have been reduced in scope where the least damage would be done by subs-

tituting information available from published sources,

Phase I of the two-phase multiyear program is underway and most  
tasks are on schedule. The first wet season sampling surveys have been  
completed and substantial progress has been made toward achieving phase

I goals. Unexpected results (e.g. the discovery that one lizard species  
which is uncommon near the ground is extremely abundant in the forest  
canopy) and elucidation of the food web by gut content analyses and feed-  
ing observations have already demonstrated the value of the phased approach.

The integration of these results with the massive body of biological inform-  
ation from the rain forest at ET Verde (Odum and Pigeon, 1970) will pro-  
vide the needed departure point for investigating energy related disturb-  
ances in this important ecosystem,

Four randomly selected plots (1 ha. each) were surveyed and marked  
in July, 1980. A 300 m transect line was subsequently placed diagonally  
through each plot. A single 300 x 300 m grid (staked at 30 m-intervals)  
for use in vertebrate surveys has recently been added,

Calorimetric measurements have been made on selected animal  
Species. Additional species will be sampled, and vegetation samples will  
be processed as phase I is continued. Sulfur content has been calculated  
for two common species of lizards and estimates for other species are

Underway. Other mineral analyses will be added in turn,

A new task has been added with the objective of quantifying carbon export from the forested Rio Espiritu Santo watershed. Data from this study can be correlated with rainfall and decomposition information ob-

---Page Break---

tained in the study area (which lies entirely within the watershed) to extend our knowledge of this potentially important process.

Vegetation studies are behind schedule as the result of an untimely staff vacancy which developed at the beginning of the field season. Research is now underway on phenology, Titterfall, and plot characterization. Decomposition studies are now scheduled to begin during the forthcoming dry season (January-April, 1981). Plot characterization and interplot comparisons are considered important in phase I so that they may serve as an accurate basis for phase II investigations.

Faunal studies have been emphasized during phase I because of the need for comprehensive data on the interrelationships among major animal groups in rain forest communities. Initial investigations have already provided significant new insights on abundance, distribution, and feeding relationships among species, and further discoveries are expected when studies are continued into the next field season.

The role of phytophagous invertebrates is central to food web analyses. Their use as prey for larger species is partially known from the literature and from initial gut contents studies, but research on their role as primary consumers has been delayed pending the hiring of an invertebrate ecologist. The position has recently been filled and studies are scheduled to begin in February, 1981.

Vertebrate studies are on schedule, with the exception of mammal surveys which were delayed because of a potential rabies problem. Key species have already been identified in each class. The tree frog *Eleutherodactylus coqui* is easily the most abundant amphibian. Three lizard species (*Anolis gundlachi*, *A. evermanni*, and *A. stratulus*) are the most abundant reptiles. Although the bananaquit (*Coereba flaveola*) is the most abundant bird species, other larger species such as the red-necked Pigeon (*Columba squamosa*) may be key species in terms of biomass. Among mammals the black rat (*Rattus rattus*) appears to be the most abundant terrestrial form. Bat species have been collected but studies of their population density are difficult and have not been attempted in phase 1.

---Page Break---

Progress at this point in phase I studied has been substantial.

Calorimetric and chemical analyses of key species are underway and carbon export research has been initiated. Vegetation studies, although started late, will produce the required data by the end of phase I field studies. Several key animal species have been identified, including one anole which was formerly thought to be uncommon in the forest. The importance of vertical studies for a1 faunal groups has been demonstrated and will be emphasized as research continues. Initial food studies have produced New information on the feeding relationships and identified some of the major pathways of movement for energy and mineral through the foodweb. Integration of information from separate tasks is inappropriate at this point, but the preliminary results suggest that new insights into animal community organization in tropical rain forests are likely to result from phase 1 studies.

---Page Break---

## 2.0. INTRODUCTION

The accelerated development of traditional and unconventional energy resources is impacting ecosystems throughout the world. The

nature and extent of these impacts is related not only to the type of energy development (e.g. fossil fuel, nuclear, biomass, solar, etc.), but also to the prevailing environmental conditions (e.g. annual tem-



erature fluctuations, rainfall elevation). Therefore information on temperate ecosystems has limited application in tropical regions. Potential impacts on temperate zone ecosystems are relatively well known but tropical ecosystems, particularly rain forests, are understood in far less detail, Their importance in the world carbon cycle and regional significance have been identified and inputs and outputs have been calculated, but the structural complexity and functional interrelationships of species (e.g. food webs, mineral cycles) are poorly known.

We are therefore investigating cycling and transport processes within a relatively simple moist tropical (tabonuco) forest. Initial studies are being conducted in the rain forest near the El Verde Field Station, Once a comprehensive overview has been obtained through integration of the results of Phase I field studies with the substantial body of published information on the same study area, in depth studies involving experimental manipulations will be undertaken.

Because the original program plan (June, 1980) was formulated prior to the reduction in funding for Fiscal year 1961 the program has been adjusted accordingly. Basic objectives and overall design have not been altered. Some task elements have been eliminated, and some sampling reduced to 2 single season where it was felt that sufficient supporting information could be obtained from existing literature, Other task

elements have been added on the basis of observations and preliminary  
?data analyses obtained during the first seasonal sampling period.

This document presents the modified program design and progress  
as of November 30, 1980 on the Phase I studies. Detailed methods are

---Page Break---

included so that this report can also serve as a reference guide for  
future studies. Extensive analyses of preliminary data are not possible  
at this point. Substantial, and in some cases unexpected results have  
been obtained which have confirmed the value of this phased approach.  
Results are presented and discussed by task in this report, but because  
of the incomplete nature of data sets no attempt has been made to in-  
tegrate data from different tasks.

---Page Break---

### 3.0 PROGRAM OVERVIEW

It is necessary to understand the major features of ecosystem

Structure and function and their response to exogenous environmental

variables in order to predict the effects of expanding energy technology on tropical ecosystems. Current knowledge of the source-sink role of tropical rain forest biota and the factors which regulate this role is insufficient to foretell the impact of energy development on the mobilization and release of critical elements or the ecosystem's capacity to assimilate elemental inputs. By investigating cycling and transport in a well studied and relatively simple tropical forest ecosystem, we intend to identify key pathways and major reservoirs of minerals and energy so that relevant hypotheses can be developed and tested concerning potential energy development impacts.

Cycling and transport processes are being investigated in the tropical moist (tabonuco) forest at E1 Verde by conducting a two phased program so that a firm data base can be established and subsequent experimental manipulations can be undertaken. The long history of environmental research in the Luguilo Mountains and the existence of the E1 Verde Field Station in this forest provide a unique opportunity for this type of research. In the proposed program we are building on previous work at E1 Verde to examine the food web structure and identify important factors involved in faunal components of cycling and transport processes, a concept which has received little attention in temperate ecosystems and even less in the tropics,

Phase I studies are focused on obtaining missing information on the forest ecosystem and integrating these data with published information

to construct a meaningful model of mineral and energy storage and movement.

Faunal components are emphasized in this phase because of the lack of

Comprehensive food web data for tropical forests, and because of the wealth

of relevant vegetation data for the study area, The objectives of phase 1

Studies are:

1. to identify the major reservoirs and pathways of minerals and energy in the forest, and

---Page Break---

2. to develop hypotheses concerning the potential effects of energy cavelopment related disturbances (e.g. inputs of sulfur, carbon, etc.) on cycling and transport.

Each task and subtask will provide information directed toward

these goals. The mineral inventory wil] provide information on the

?mineral and energy value of each major class of items (Fruit, leaves,

feces, individual animals, etc.) in each ecosystem compartment. Faunal

studies will quantify the number of such items in each compartment. and

the rate of movenent of these ?tens between compartments, In addition

to studies characterizing study plots, the vegetation work will provide

@ basis to compare the magnitude of elemental flow through compartments with direct movement of primary production to the decomposer compartment via leaf and fruit fall and litter decomposition.

We are attempting to achieve the above-stated objectives by the following steps:

1) delineation of trophic structure food webs and food web segments,

2) identification of dominant species in each food web or food web segment,

3) quantification of vertical, horizontal and temporal pattern of the distribution of identifiable functional units (species, trophic groups, food web segments, etc.) of the forest ecosystem,

4) quantification of the distribution of key elements among major functional units of the system taking into account Spatial and temporal considerations

5) development of a refined forest ecosystem model which will Permit simulation of cycling and transport processes, and

6) generation of hypotheses concerning the potential impacts

Of energy development which are both relevant to tropical forest ecosystems and testable in phase II,

An important result of the proposed study will be a detailed food web in which major aspects of the distribution and transfer of minerals are known. Evaluation of these data is likely to produce useful insights on the relationship of food web complexity to species diversity, and about the influence of rainfall, humidity, soil type,

---Page Break---

nutrient pools, dominant consumers, and food web structure on the overall structure of ecosystems. This information will provide an important conceptual basis for inferring the key points at which perturbations due to energy development are most likely to disrupt natural systems.

Phase II will primarily involve the testing of hypotheses, although

Some phase I studies (e.g. population turnover rates) will be continued into this phase. Hypothesis testing may take a variety of forms. Direct manipulation of the forest by methods such as the addition of particular elements may be undertaken, as permitted by the U.S. Forest Service.

Studies of existing plantations of various ages and managed forest areas may be conducted to gather comparative data on cycling and transport

characteristics, including food web complexity. The exact nature and extent of these studies will depend largely on information acquired and synthesized in phase I and are not considered in detail at this time, although Several questions relating to the impacts of energy development on tropical forest can already be posed.

Several hypotheses can be generated which involve comparisons between native tropical hardwood (tabonuco) forest and plantations (including those which are managed for energy production). Specifically, one can hypothesize that in the native hardwood fores:

1. productivity is higher,
2. consumer biomass is greater,
3. insect pests are less abundant,
4. food web structure is more complex,
5. nutrient cycling is more rapid, and
6. the system is more resilient to exogenous disturbances

than in plantations on the same soil types and under the same rainfall

regimes. The hypotheses selected for testing in phase II will be based on information acquired in phase I and on a realistic evaluation of potential energy related impacts.

---Page Break---

#### 4.0 PHASE 1 METHODS

A compartmentalized ecosystem model (Figure 1) is being used as a framework for coordinating the various task studies. York will focus on those ecosystem compartments above primary producer in the food chain for the following reasons: (1) the importance of higher trophic level in mineral transport has already been suggested for some forests (Weir, 1969), but the role of fauna in moving energy and nutrients among compartments of terrestrial ecosystems has been largely neglected (Sturges et. al., 1974; Burton and Likens, 1975), and (2) primary production and elemental cycles have already been studied in the moist tropical forest (Odum and Pigeon, 1970). For these reasons only a partial measurement of mineral transport in and out of the forest ecosystems will be begun in phase I of the project. and values needed for the current study will be taken from the literature and/or from the work of visiting and collaborating scientists.



## 4.1 Sampling Design

The study area (Figure 2) was selected because of its long history of continuous research, beginning with the rain forest gamma radiation Studies (Odun & Pigeon, 1970), and its proximity to the established facilities at the £1 Verde Station. Other factors which were evaluated in

the study area delineation process were the relationship to U.S. Forest Service research areas and the amount of existing disturbance from previous studies.

The overall design is stratified random, Four sampling locations (Points) were randomly selected within the study area so that subsequent Statistically valid analyses can be performed (Green, 1979). Sampling points were selected using a grid technique (Phillips, 1959). The following criteria were used in the selection of random plots:

© potential confounding factors (e.g. roads, perennial Streams, previous destructive sampling) are not present within 50 m of a sampling point.

---Page Break---

Figure t, Generalized Ecosystem Model

10

---Page Break---

Figure 2.

SCALE IN METERS

Rain Forest

Study

Area

---Page Break---

© sample points will be no closer than 200 m to each

?other so that sampling overlap can be avoided, and

© reasonable access is possible.

?The size of the area to be sampled at each randomly chosen

Point will vary according to the discipline being studied. A one hec=

tare plot with boundaries marked at 10 m intervals has been established

at each sampling point. Plot and plotless methods, transects, and detail-

?ed inventory procedures are used in phase 1 investigations. Studies

focus on sampling points, but include other portions of the defined study

area,

Most sampling 1s being conducted during a minimum of two seasons.

Most surveys will occur during the wet and dry seasons, but additional

Sampling 1s required for some tasks (e.g. breeding bird surveys). The

timing of field surveys is being coordinated to avoid sampling interference yet maintaining close temporal correspondence among field studies.

## 4.2 Task Methods

Each task is described as a separate entity within the overall phase I sampling design because of the variation in pertinent information already available for different disciplines. The tasks are of varying durations, but all should be completed within 12 to 14 months (see Schedule section). Following the initial phase, the program will be evaluated and some studies continued with modifications (e.g. growth and population turnover of key species). Although the data acquired should be of immediate value, they are also intended to provide a basis for long term in depth studies continued under phase II of the cycling and transport program. The options will be better defined as phase I approaches completion,

Objectives, detailed methods and results are presented for each task and the significance discussed. Conclusions and integrated data presentation are not included because of the preliminary nature of the information obtained for some of the tasks.

a1.

---Page Break---

### 4.2.1 Physical Phenomena

Basic information on physical aspects of the environment is necessary in order to interpret biological observations and to provide baseline data for phase II studies. Specific objectives are

# to collect pertinent weather data to be used in evaluating information collected from other tasks

© to document seasonal and other temporal changes within the study area

# to record stream flow from the Sonadora watershed in order to evaluate transport from the study area

## Materials

# rain gages (at £1 Verde Station and tower)

?© thermometers and hygrothermographs (at £1 Verde Station and four Towers at the tower)

wind speed and direction recorder (tower only)

stream gaging station (at suspension bridges on Quebrada Sonadora within the program study area)

#### 4.2.2 Mineral Inventory and Energy Studies

The fixation of energy from the sun in plant materials is the ultimate driving force for cycling and transport processes in tropical rain forests. Movement of key minerals and energy through the systems

is being investigated by identifying the major pathways and quantifying movement between major compartments (Figure 1). Because of the important role of tropical forests in carbon cycling, a new task has been added which addresses the role of rainfall events in carbon export from the watershed which includes the program study area.

##### 4.2.2.1 Mineral Inventory

Sulfur: Material to be analyzed for sulfur is mixed 4:1 with benzoic acid, compressed and oxidized by burning in a combustion bomb under 28 atmospheres of oxygen pressure. The contents of the bomb are subsequently depressurized and rinsed through a coarse, rapid filter with four

air

---Page Break---

50 ml. portions of doubly distilled water. The volume is reduced to 100 ml. by evaporation on a hot plate and standard gravimetric techniques are then used to analyze for sulfate,

#### Other Methods

All other analyses are carried out according to procedures specified in APKA-AMNWA-HPCF "Standard Methods for the Examination of Water and Wastewater", nitrate using the standard amalgamated cadmium reduction followed by diazo coupling of sulfanilamide and N-(1-naphthyl-ethylenediamine). The intensity of color is read with a spectrophotometer,

Organic and ammonia nitrogen are measured using standard Kjeldahl analysis with titration of indicator boric acid solution following digestion and distillation.

Phosphate is measured using vanado molybdate heteropoly acid color development and subsequent readout using a spectrophotometer.

Collections associated with the sediment graph/carbon export correlation are also being used for analysis of export of nitrate, phosphate and organic nitrogen. The possibility of sulfur/sulfate analysis will also be addressed in the coming months.

Inasmuch as wet weight/dry weight and ash weight analysis of

secondary consumers such as eleutherodactylid and anolid species in the forest are not well described, and since ash-free dry weight is important for the description of the calorimetry of these animals, data on ash weight, wet and dry weight are presently being accumulated. The animal's tissue is 4150 burned under controlled conditions (bomb calorimeter) in order to determine the energy storage in the animal compartments of the forest.

The residues after burning were rinsed, filtered and analyzed for sulfur (30,7).

#### 4.2.2.2 Calorimetry

Calorimetric measurements are being made on organisms which are known or suspected of being major components of the rain forest ecosystem.

a1.

---Page Break---

Dried samples are mixed with benzoic acid burn adjuvant and an accurately weighed sample of approximately 1 gram is burned in the oxygen bomb at 28 atmospheres pressure following moistening with 2 drops of a 1% solution of triton X-100, These conditions control the burn, Standards of pure benzoic acid (6,318 cal./gn) are run periodically as a calibration of the calorimeter, Temperature differences are read to the nearest thousandth of a degree. All measurements are corrected for unburned fuse wire and a pre-burn oxygen purge is used to eliminate atmospheric ni-



trogen from the borb,

#### 4.2.2.3 Carbon Transport

This task was added subsequent to the development of the original Program plan. Carbon transport from the forested upper Rio Espiritu Santo watershed is being quantified, Analysis is based upon a model which wages sediment export from flow-monitored streams. The Cara del Indio hydrological gaging station on Rte. 966 is an appropriate site for collections of sediments for analysis of the sediment graph during Sediment exporting events, The hypotheses developed are based on the following considerations:

1. Sediment may or may not be associated with the bulk of the carbon exported via the drainage system.
2. Base flow conditions which might be responsible for Some inorganic (dissolved) carbon export should not be very significant,
3. Long term storage cycles of carbon ( > 1000 years) may be responsible for more CO<sub>2</sub> atmospheric buffering than is presently thought.

Our observations consist of surface and bottom water col-

tection during base flow and high water conditions and analysis of the water with respect to inorganic and organic carbon.

Organic carbon is measured using a standard sealed vial oxidation technique (c.f. Oceanography International Operating Manual) employing potassium persulfate as the strong oxidant. The vials are scored and broken after the neck has been sealed within a gas carrying system. A

a5.

---Page Break---

urge of nitrogen gas acts as a carrier to bring the liberated carbon dioxide (CO<sub>2</sub>) through the infrared sensing system of an Oceanography International total organic carbon analyzer. The detection response is integrated with respect to time electronically and the relative integrated peak area is related to calibration curves. Substances used for calibration of the response curve areas include sodium carbonate primary standard and potassium acid phosphate primary standard,

Inorganic carbon is measured by injecting a known volume of sample into a 2M solution of phosphoric acid placed within the nitrogen purge system of the Oceanography International total carbon analyzer. These acidic conditions are sufficient to convert all carbonic acid, carbonate and bicarbonate ion to free carbon dioxide which is purged with the nitrogen system described above,

### 4.2.3 Vegetation Studies

Considerable information has been accumulated on the vegetation of the Luquitlo Mountains, including the rain forest at El Verde (Odun and Pigeon, 1970). Phase I studies are designed to collect important data on 597s (e.g. variability in Titterfall, decomposition rates, etc.) which have been identified, and will provide baseline data for experimental studies which are planned for phase II.

#### 4.2.3.1 Plot Characterization

Several types of information are needed in order to provide a basis for interpreting faunal data and for designing phase II investigations. Initial characterization techniques are described below.

#### Structural Analysis

All trees with a diameter at breast height (DBH) of 10 cm or greater will be identified, tagged with species name, DBH, measured, and height determined,

Between plot comparisons will then be made of:

Species composition,

Species importance,

-16=

---Page Break---

Biomass, and standing crop of elements in biomass

(Ovington & Olson, 1970).

Topography

Information will be gathered on slope and aspect in each site to add information regarding between-plot variation.

Loose Litter

# Variations in the forest floor litter components among plots will be determined by collecting samples at random in each plot each season. The number and size of the samples will be based on plot heterogeneity and prior sampling surveys.

\* The material collected will be separated into the wood

and miscellaneous components, dried at 70°C for 72 hours, and weighed

Pilot studies were conducted within the study area in April, 1980 which indicated that 1 hectare was adequate for phase I vegetation studies. Further work is not planned in this phase,

#### 4.2.3.2 Phenology

##### Objectives:

- \* To look for temporal differences in species flowering and fruiting patterns within and between plots

- \* To determine importance of species contribution to Flower and fruit fall

##### Methods and Materials:

- \* See Titterfall for sampling location information

® Flowers and fruits will be separated monthly by Species for each basket, counted, dried at 70% for 72 hrs. and weighed.

- \* Phenological studies will be limited to the Flowers and fruit contribution to litterfall as determined by basket collection in each plot. Prior extensive Phenological studies in P.R. (Estrada, 1970; Bannister,

?We

---Page Break---

1970, Nevling, 1971) indicate that detasted ad-  
ditional studies would be repititious and of low  
Priority at this time,

#### 4.2.3.3 Litterfalt

Objectives:

to determine seasonal pattern of litterfall as

@ whole and for individual species

© to detect dfferences between sites for total

and species Htterfall

Nethods and Materials:

# Twenty ten? galvanized hardware cloth baskets lined with 1 mm mesh fiberglass screen were placed in each of the 4 plots.

© The Locations of 4 Titterfall transects were randomly selected at each site with 10m intervals between potential transect lines. Five baskets were randomly located along each? transect (Figure 3).

# Each basket was elevated at least 10 cm from the forest floor (30 cm where possible) and leveled.

\* Collections of the Titterfall are made bimonthly. The mid-month collection is separated by basket into leaves, flowers, fruits, wood and miscellaneous items,

© The collections at the end of the month are air dried; separated by species into leaves, flowers, fruits; and sorted by basket into wood and miscellaneous items.

© Each component at each collection is counted, dried at 70°C for 72 hours and weighed.

Her season-dry season samples will be processed by

species by plot (some species by transect) for: cator=  
ic and mineral content.

#### 4.2.2.4 Leaf Decomposition

Objectives:

\* To determine the seasonal decomposition of freshly fal=

Ven Teaves

© To compare the between plot rates of leaf decomposition

1

---Page Break---

Figure 3.



Randomly Selected Vegetation Sampling Locations Within Study Plot.

-19.

---Page Break---

© To evaluate the effect of species composition of the decomposition rate

© To determine the change in caloric value and mineral content over time and between major seasons.

## Methods and Material

© Freshly fallen leaves will be collected seasonally + from each of the four study plots in order to represent species composition, approximate quantity, and the chemical composition of the leaves in each plot.

The leaves will be air dried overnight to remove surface moisture. Twenty grams of air dried leaves will be placed in each 25x40 cm fiberglass screen

bag (Imm mesh) for a total of 170 bags/site,

© Eighty bags containing leaves from plot 1 will be placed in plot 1, four near each of the 20 litter baskets. Eighty bags containing leaves from plot 1 will be placed in plot x selected for site homogeneity. Ten bags will be reserved for fresh weight-dry weight conversion.

# The above will be conducted for each of the four plots, thereby detecting any variation in the decomposition rates between plots. By randomly placing decomposition bags from each of the 4 plots in one homogeneous area (plot X), any differences in the decomposition rates, between the plots due to species composition will be determined,

© Ten bags will be collected at random from each site plus 10 bags representing each site from plot X (40 bags total each collection) at 7, 14, 20 days; 2, 4, 8, 16 months. The remaining bags will be considered a field reserve in case of unpredictable bag destruction.

\* Caloric values and mineral content will be determined on 3 of the bags from each plot at each col-

lection to determine temporal variations.

#### 4.2.4 Faunal Studies

The roles of animals in ecosystem structure and function can be effectively evaluated by combining energy flow and nutrient cycling studies. Recent work in temperate forest ecosystems has provided valuable information on the roles of animal populations as nutrient "sinks" and as agents of nutrient transport (Weir, 1969; Sturges et al., 197

20

---Page Break---

Because of their potentially high turnover rates at relatively high ambient temperatures, underestimating the importance of organisms with short life spans and low standing crops in cycling and transport processes is even more likely in tropical systems than in temperate systems. The proposed phase I faunal studies are designed to investigate that possibility in the tropical forest ecosystem at El Verde.

The major faunal sampling effort is aimed at obtaining inform

attention on the food habits and general abundance of animals. From these data emphasizing top carnivores and herbivores a comprehensive food web will be constructed. Initial food web analyses will attempt to define major pathways rather than focus on parallel food subwebs (Paine, 196: Gilbert, 1980) which may or may not be important structural units in this insular tropical rain forest.

Studies will focus on those intermediate consumers serving as Principal diet items for higher order carnivores. Attempts will be made to estimate population turnover rates for key (i.e. abundant, frequently eaten or Target) consumers. To some degree, intermediate consumer roles and rates will have to be estimated in a pooled fashion from differences between rates of primary productivity Vitter fall and decomposition and Consumption rates of top carnivores.

The faunal studies will phase from highly diversified qualitative collection through more selective sampling aimed at relative abundance to highly specialized quantitative sampling, aimed at key species. Relative abundance and absolute abundance data will be combined to refine portions Of the food web. Turnover estimates plus metabolic data will be used to

quantify flows. Because of expected seasonality some types of sampling

WITT be continued throughout the entire period of study. However, with

each successive sampling period, more effort will be placed on relative

and absolute abundance estimation for key species and species groups rather

?than upon the identification of new species.

In time, more detailed paths will be able to be delineated and

quantified and it will become possible to test whether additional re

finement is needed to predict the influence of man-made changes upon

<2.

---Page Break---

Processing by these ecosystems. More specific descriptions of research tasks and methods follow for the major faunal groups.

#### 4.2.4.1 Invertebrates

?The invertebrates fauna influences cycling and transport proces-

Ses in the rain forest ecosystem in three major ways. First, by grazing

for photosynthetic and reproductive structures of plants (e.g. phytophag-

Sous insects) they affect forest primary productivity and alter plant

Species composition and diversity (Janzen, 1970, 1973; Mattson and Addy, 1975). Secondly, as decomposers and detritus feeders in the soil and litter they are responsible for breaking down dead organic matter, making nutrients available to the plant community. This latter function is of great importance to the plants of a tropical forest, which, due to thinning and soils low in nutrient concentrations, must incorporate nutrients at a rapid rate. Thirdly, invertebrates play a role of unknown magnitude as predators, certainly influencing and possibly controlling the density and species composition of selected primary consumers (Hairston et al., 1960).

In view of these characteristics of the invertebrate fauna, the importance of their role in the flow of energy and cycling of nutrients in the tropical rain forest becomes evident. Unfortunately, few studies have contributed to the understanding of the different ways in which invertebrates influence the flow of energy and nutrient cycling in tropical terrestrial ecosystems. The majority of invertebrate studies in the tropics have been limited to crude estimates of absolute or seasonal abundance and some biomass determinations.

Studies of the dynamics of invertebrate populations have historically emphasized economic pest species, resulting in models of populations that are not general enough for prediction of randomly selected species (Clark, et al., 1967; Gilbert and Singer, 1973). The few detailed studies of invertebrate populations in the tropics, limited to groups of insects such as butterflies of the genera *Acraea* (Owen, 1971) and *Heliconius*

(Ehrlich and Gilbert, 1973), and euglossine bees (Janzen, 1971), have shown

-22-

---Page Break---

that for these species population size remains constant through time

but turnover rates are high. If this is true for other tropical forest invertebrates, then because they comprise a large proportion of the faunal biomass, invertebrates must play a major role in tropical ecosystem nutrient cycling and energy flow. It has been demonstrated in temperate forests that invertebrates, in particular phytophagous insects, regulate forest primary productivity (Mattson and Addy, 1975). However, the same has not yet been demonstrated in tropical ecosystems.

Most of the work concerning forest invertebrates in Puerto Rico has been limited to pest species, taxonomic descriptions, and relative abundance and/or food preferences of selected species (Martorell, 1943; De León, 1944; WoTeatt, 1940; 1986; 1950; Drewry, 1970). The only studies that have emphasized the role of invertebrates in the rain forest ecosystems were performed during the AEC Radiation Experiment (Odum and Pigeon, 1970) in the tabonuco forest at El Verde. These studies covered the major groups of invertebrates in the tabonuco forest including aquatic communities in bromeliad leaf axils (MacGuire, 1970), nematodes in litter and soil (Coleman, 1970), termites (MacMahan, 1970), soil microarthropods (MacMahan &

Solins, 1970) and mosquitoes (Weinbren & Weindren, 1970). However, these Studies were fragmentary and conducted during different seasons and/or years and leave even the basic inventory incomplete.

To attempt to study the role of all invertebrates in ecosystem Structure and function and their effect on energy flow and nutrient cycling {5 impossible due to the magnitude of the task. Additionally the ?taxonomy, autecology and population dynamics of most invertebrates have not been well studied, For this reason the scope of this study will be limited to phytophagous insects, their principal predators and the soil fauna.

© Phytophagous insects: Phytophagous insects of both Tow and high strata of the vegetation will be sampled by means of ?Sweep nets special enclosures, malaise traps, light traps and/or a sucking apparatus (De-Vac Sampler). Sampling wit! be done at each season both day and night. Environmental variables will be recorded during sampling.

23

---Page Break---

?The effect of phytophagous insects on energy flow and nutrient cycling will be evaluated in coordination with related vegetation studies. The intensity of grazing on Teaves and fruits will be quantified by means of the relation



ship between leaf area damaged or consumed and leaf area index through time.

Soil fauna: Approximately 90% of forest primary productivity goes to decomposers (Gray, 1964; Wiegert, 1970); therefore they will be considered the most important with respect to nutrient cycling and energy transfer. The role of soil fauna will be approached in the same manner as for phytophagous insects,

Because sampling accuracy is of utmost importance in estimating population size, soil litter fauna will be sampled by more than one technique. Sampling techniques will be tailored to the characteristics of the different groups. With the help of expert consultants the best method will be chosen from the literature. These techniques may include soil coring (nematodes), flotation, extraction (Berlese and Tullgren funnels or high gradient canisters for microarthropods). Direct captures will also be utilized for macroarthropods such as arachnida. As with the phytophagous insects, dominant species will be identified and future experiments designed to investigate their population dynamics.

The soil and litter fauna will initially be treated as one compartment with only prey species being singled out as they are identified from studies of carnivore food habits. As

Stated previously, for most samples only information on total numbers or biomass will be secured, However, attempts will be made to determine the relationship at different seasons of the year between numbers of cores (or area sampled) and numbers of taxa recognizable,

24

---Page Break---

#### 4.2.4.2. Amphibians and Reptiles

These poikilothermic vertebrates are conspicuous and abundant components of tropical forests. Considerable data are available on the ecology of some groups (e.g. the genus *Anolis*), but the kind of information required to determine their importance in cycling and transport through these ecosystems is incompletely known. The basic type of data required are (1) mineral and energy content, (2) status in food web (Food habits and predators), (3) biomass, and (4) population turnover rates. Initial studies will focus on acquiring information of the first three types.

##### Subtask 1. Species Inventory

Objectives:

# to determine the presence and general abundance of species

not readily observed by transect methods

## Methods and Materials:

\* Intensive searches at random and selected locations will be conducted periodically throughout the study area, and random observations will be noted.

© Surface debris and litter will be overturned in order to locate secretive and fossorial species.

# The species, location, and general abundance of each species observed will be recorded.

## Subtask 2. Presence and Relative Abundance

objectives:

\* to determine the species composition and relative abundance of amphibian (anuran) species within the study area in order to identify important species and to correlate information collected at study plots with more detailed data collected on coguis within the same general area by Dr. M. Stewart and her colleagues.

## Methods and Materials:

© Conduct transect surveys along same lines as bird and Vizard surveys,

© Walk each transect on three separate evenings at each major season (wet and dry) and record the species, sex

25

---Page Break---

(where possible), and size class of each individual observed along a 50 x 2m belt transects

© Record the calls of any individuals which cannot be identified in the field,

© Calculate the relative abundance for each species in each of the four plots.

### Subtask 3, Anolis Food Habits

#### Objectiv

© to determine the types of food taken, frequency of occurrence, and percentages of food items for each species,

Methods and Materials:

© Collect a minimum of 30 adults of *Anolis gundlachi* and *A. vermanni*. during each major season (wet and dry)

© Skin, weigh, and cool specimens immediately in the field,

© Dissect stomach content out of each specimen and place them in preservative (10% formalin or commercial insect preservative). General references Sexton (19/2).

© Identify contents as specifically as possible, noting approximate percentage of total contents for each individual lizard,

#### Subtask 4. *Anolis* Population Density

objectives:

© obtain a minimum population density estimate for common species of *Anolis* in each plot at each season (wet & dry)

© obtain a population index and relative abundance estimate for *Anolis* in each plot.

Methods and Materials:

© A permanent transect 150m in length beginning at the center of the plot and extending diagonally from it along existing

bird transect is established in each of the four study plots.

© Each transect is surveyed by slowly walking the marked line on three separate occasions: morning (0700-1000), mid-day (1000-1400) and afternoon (1400-1800) during the wet and dry seasons,

# The species, sex (or size class), and distance from the

centerline of the transect is recorded for each individual observed.

=26-

---Page Break---

Relative abundances are calculated for each plot, and minimum population densities are calculated according to Frye's strip census formula (Overton, 1971):

$\bar{F}_i$ : average perpendicular distance

$B_i$  between observed animal and transect centerline

$C_r$  correction factor for units of measurement

transect length

area of study site

estimated population density

number of animals observed on

transect

#### Subtask 5. Anolis Population Turnover and Growth Rates

##### Objective

© to obtain reliable estimates of population turnover and

Growth rates for the most abundant species of Anolis

##### Methods and Nateria:

© Live capture all sizes of Anolis within a designated

Portion of the overall study area.

# Mark animals individually, weigh, measure, sex (when

Possible), and release. Repeat at 1-3 month intervals  
and extend into phase II.

4.20

### 3 Birds

Synthesis of new techniques and knowledge gained from preliminary

Field work has resulted in some changes in the original work plan, with a  
view toward increasing the value of the resulting data. Current methodology  
for the program is described below.

#### Subtask 1. Population Density

##### Objectiv

© To obtain reliable population estimates for each species  
in all sites studied and to detect changes in population  
density throughout the year.

27

---Page Break---



## Methods and Material:

© Grid 9 ha. area that includes one of the sampling plots (Plot 3). Grid lines will be established at 30m intervals and intersections marked with numbered Fiberglass stakes,

© Beginning in February, 1981, use modified spot-map methods, mist netting and systematic observations, to determine population density for each species in the 9 ha. site (see Holmes and Sturges 1975).

© In each of four 1 ha, sampling plots, establish a 300 m transect line through opposite corners of the Plot. Set up a 150 m net lane bisecting each transect line,

© Visit each site 2 days/month to conduct two transect Censuses between 0600-0900 using the method of Emlen (1971). Every third month run four transects in each site between 1500-1800.

© By comparing absolute densities from Plot 3 with data {from transect surveys from the same plot, calibrate

the transect densities to the number of birds actually occurring, ?Use transect data to estimate population densities in the other three sites.

© After initial transect counts, sample each site dawn to dusk for five consecutive days with mist nets, marking a) birds captured. Calculate trappable population sizes (MacArthur and MacArthur, 1974).

## Subtask 2. Feeding Behavior

### Objective

© To accumulate sufficient observations of feeding to test for differences between species, sites, seasons and individuals.

### Methods and Material

\* Observe avian behavior for equivalent periods in each Site. Record prey items taken, horizontal coordinates Of prey capture, height of prey capture, tree species, type of foraging move, direction and distance of foraging move, substrate attacked, perch diameter, location, time, weather conditions, etc.

© Data will be compiled and analyzed in a fashion that will make possible comparisons with foraging data taken by C. and E. Kepler in Colorado forest and A. Cruz in pine plantations.

---Page Break---

Subtask 3. Diet and Weight

Objective:

Methods and Materials:

To determine differences in diet and weight between species, sites, seasons and individuals; to describe food webs for each site

Sacrifice sixty birds of 10 species captured outside study sites, identify stomach contents and freeze for bomb calorimetry.

In conjunction with the banding studies in Plot 3, use ground and canopy mist nets to capture as many individuals as possible in this plot. Also do occasional mist

netting around the boundaries of the other three plots,

Weigh, measure and mark each individual capture,

Sample stomach contents of each individual with antimony  
potassium tartarate enetic.

Identity stomach contents to the lowest taxonomic group  
possible.

Construct partial food web for each site using data col=  
Tected and literature sources

Calculate uptake of minerals using feeding rate and  
bonb calorinetry data,

Subtask 4. Materials Discharge

Objective

Methods and Materials:

to obtain reliable estimates of the amount of fecal  
and regurgitated material produced/individual of each  
species per unit tine

Place mist-netted birds in holding cages for 1) one hour during the day and 2) overnight.

Collect, dry and weigh defecated and regurgitated material.

Determine average weight of feces and regurgitated material

and Bond fecal and regurgitated material to determine mineral content.

Using observed rates of defecation in the wild and in birds caged overnight, calculate amount of material returned to environment/individual/unit time,

2290

---Page Break---

@ Test for differences between individuals, species and sites,

Subtask 5. Mineral Content

Objective

© To obtain reliable estimations of the mineral content per unit weight of 1) whole stomach contents and individual food items, 2) body tissue, 3) feathers and 4) feces and regurgitated material.

#### Methods and Materials:

© Capture and sacrifice sixty birds of ten different species.

@ Remove stomach contents, identify and count, dry separately. Remove and combine fecal material from each species and dry.

# Remove feathers, and homogenize and dry feathers and body tissue separately,

© Determine important food items for each species and collect, homogenize and dry samples of each item.

# Bond and analyze residue of samples in the following numbers: body tissue, up to 6/species or until individual variation is established (minimum 10, maximum 60); feathers, same as body tissue; stomach contents, 6/species (total 60); fecal material, 2/species (total 20); food items, up to 50. Grand total? = 150-250 samples.

© Calculate per item or per event transfer of minerals.

#### 4.2.4.4 Mammals

?The role of mammals in the rain forest ecosystems of Puerto Rico 1s

Of particular interest because of the low species richness and because the larger species (e.g. black rat and Indian mongoose) have been introduced by

?man. Studies of this group will be limited in scope, but basic information

1s needed prior to implementation of experinental investigations which are plan=  
ned for phase 11,

Ojectiv

?# to determine the presence and general abundance of  
species within the study

30

---Page Break---

Methods and Materials:

© Twenty Sherman Tine traps (3x3x9 in) are placed in two lines of 10 traps each at 15m intervals between lines and traps and checked each day for a minimum of three consecutive days. The traps are checked and rebaited (as necessary) daily. Lines were surveyed during mid-November, 1980 and will be surveyed again during the 1981 dry season.

© Larger terrestrial mammals are sampled using Tomahawk line traps (6x6x24 in) deployed in a 5x5 trap grid (Nellis, personal communication) with traps spaced at 30m intervals. Traps are checked and rebaited daily for a minimum of three consecutive days.

© Bats, the only native mammals presently known to inhabit the Study area, are surveyed using horizontal and vertical mist nets erected in selected locations within the study area at intermittent time intervals throughout each major season.

31

---Page Break---

## 5.0 RESULTS AND DISCUSSION

Initial surveys have produced substantial, and in some cases unexpected results. Because these data are preliminary, integrative analyses are not possible at this time, Results are herein reported



and discussed by task,

## 5.1 Physical Phenomena

Daily rainfall and continuous records of relative humidity and temperature fluctuations have been collected at the El Verde Field Station since the radiation studies in the rain forest during the mid-1960's. The tower station was established in July, 1980 and will be maintained throughout the program field studies. Prior to the implementation of phase II studies a wind (velocity & direction) gage, rainfall event recorder and stream flow gage may be added,

## 5.2 Mineral Inventory and Energy

### 5.2.1 Mineral Inventory

A limited number of samples have been analyzed for percent sulfur content. Preliminary results on three species of lizards (*Anolis gundlachi*, *A. evermanni*, and *A. stratulus*) indicate similar values (0.2-0.3%) for all species. Additional individuals will be analyzed and other focal species processed during the remainder of phase 1.

Contents of the bomb calorimeter were subjected to sulfate analysis in order to determine the total sulfur content of the samples turned,

Much of the experimental work performed was dedicated to testing methodology which appeared to be just adequate for the sulfur levels encountered in the animal specimens, The results and the estimated uncertainty in results prompts us to review methodology and begin examining other alternatives for  $\delta$  analysis; this is particularly necessary for soil sample examination. Results are estimated to be uncertain by as much as 50% in the worst case and the probably uncertainty in other values is probably at least 20%,

232.

---Page Break---

Table 1 presents averages values of the analyses for organic

nitrogen (total Kjeldahl nitrogen), nitrate ( $\text{NO}_3^-$ ), and nitrite ( $\text{NO}_2^-$ )

in 8 samples collected from the surface and bottom of the Rio Espirity

Santo Confluence at Cara del Indio, The organic nitrogen concentration

Fluctuations appear to be related to those observed for organic carbon,

the highest value reported when the highest observed level of the stream occurred,

### 5.2.2 Calorimetry

Wet weight vs dry weight results for the anolid Tizards Anolis

Gundlach and *Anolis evermanni* are depicted in Figure 4, It is clear that individuals whether male or female all retain about the same amount of labile moisture. By dry weight is meant the weight of the animal after 36 hours of drying in a convection oven at 80°C.

The relationship of wet weight to dry weight for *Eleutherodactylus coqui* is shown in Figure 5. The individuals of *E. coqui* apparently retain substantially more labile moisture than the individuals of the anolid species. Approximately 81% of the *E. coqui* was labile moisture whereas the anolid species contained about 76% moisture by weight.

Heat content of anolid lizard species was determined using bomb calorimetry, The results are tabulated in Appendix A on an "ash-free dry weight" basis, The average heat content for all anolids was:

4322.1 cal/ash-free gm. with a standard deviation of:

315.4 cal/ash-free gm.

### 5.2.3 Carbon Transport

Work has progressed on a new task, the quantification of carbon transport from the forested Rio Espiritu Santo watershed which

includes the program study area, Carbon export from the forest is based  
Upon the model for the hydrograph of the Rio Espiritu Santo River present=  
ly under development by a visiting scientist, Or. Oswaldo Rendén Herrero,  
That model gages sediment export and weathering using hourly gaging and  
Sediment collection from a flow-monitored stream having @ cross-sectional  
area calibration,

ae

---Page Break---

#00"0 + L00°0 e?0°0 + 120°0 s0°0 + us"0 St'0 # 82°09 Sz 4aquenoy

£000 + 900°0 490°0 # oLt"o iso F sso ?2°0 F 6b°0 LL saquanon,

200°0 ¥ 910°0 900°0 + g10°o 070 + 06°0 62°70 F surt Of 4990329

200°0 + 00°0 400"0 + o10°0 toro + e1°o 80° + l2"0 6 4990299

woHOE wee WHET ?wep any Tose

(1/06) org

Wor sesauaou0g 931.431N/93249 IN

wolaesquaau0g uabouziN 1uePLafy 1e20)

?wosReRS Sub [dues o¥pu [ap ee] Wo.y san{eA az1L431N puE \*3zRIDIN fUuabOWIIN ?1 aLqeL

ate

---Page Break---

en toreg )

Figure 4, Wet Weight/Dry Weight. Rel.

and

relationships for Anolis

vermann

?gundlachi

---Page Break---

VO ZO

7

(wien Ka) cogs a

Bip imbos -F 40j diysuoyo

9 THAIDOPOIAUINETT 40 d1ysuojsojay

20

sW6 uy JuBIEM Kad

<0 ro £0

T

ee 4UBIOM Asa 7 AyblOm,

on

ANB 19M 410 /1NB19M FAK ?G aunbis

---Page Break---

We hypothesize that long term storage cycles are of limited importance in tropical rain forest. The long term cycle has a requirement for humus and organic soil formation which while important to temperate zone forest carbon fixation is probably of limited significance in the tropical forest. The temperature and humidity in the humid tropics are responsible for rapid decay of leaf material and the 3-4 heavy sediment exporting rainfall events effectively scour parts of the forest ground surface and the sub-surface parts of the forest ground surface and the sub-surface soil layer of its potentially fixable carbon.

Stream Chemistry and Gaging: Water samples were collected from the surface and bottom of the Rio Espiritu Santo River confluence known as Cara del Indio (Indian Face). Samples were processed both before and after either filtration or centrifugation to test the relationship between sediment export and nutrient loss from the forested watershed.

Appendix B lists data for 5 surface and 5 bottom samples collected on each date. The gaging depths are in USGS units and were taken directly from the depth marker at the USGS gaging station at Cara del Indio on Rte, 966. The designations: F and U refer to filtered and unfiltered samples respectively.



Table 2 shows the average values for carbon analysis of § samples collected on each of the 4 dates given. The inorganic carbon analysis Standard deviations indicate that the collection procedure is adequate fo ensure homogenous bulk water samples. Unusually high values of organic carbon observed for bottom samples collected October 9, 1980 and Nov. 25, 1980 may be artifacts of the sampling procedure, and, at least in the case Of the November 25, 1980 sampling a larger number of samples could have Provided statistically valid basis for rejection of the high value.

On November 17, 1980, the stream level at Cara del Indio was near base ?low condition following a period of very heavy showers during most of the previous 12 hour period. The flushing of labfle or available carbon from the watershed may account for the Tow concentration of dissolved organic carbon observed for the 5 samples collected on that date. The only data supporting the hypothesis that heavy export of carbon 1s Targe=

-37-

---Page Break---

Table 2. Carbon Concentrations of Samples from Cara del Indio.

eee

Surface

Date Organic Inorganic Flow Level

Filtered Unfiltered Filtered Unfiltered (USGS Units)

Oct. 9/80 3.0071.19 3,7941.42 5.5541.28 6.5841.21 2.82

Oct. 30/80 5.3130.49 8,380.45 3.6340.13 3.5440.06 4.46

Nov. 17/80 0.5840.13 0,8640.21 5.6840.20 5.4440.98, 2.70

Nov. 25/80 4.3241.88 4.0542.41 - - 3.50

Bottom

Date Organic Inorganic Flow Level

Filtered Unfiltered Filtered Unfiltered (USGS Units)

Oct, 9/80 5.5643.79 3,354.76 5.7241.48 5,8841.24 2.82

Oct. 30/80 5.4840.17 8.6440.57 3.6040.22 3,670.16 4.46

Nov. 17/80 0.4540.09 0.5740.16 5.7240.37 5.0840.87 2.70

Nov. 25/80 3.9042.84 4.0043.70 - - 3.50

eee

---Page Break---

ly associated with sediment exportation derives from measurements

?taken October 30, 1980 when the stream level was the highest of any

of the collection times. The differences in carbon concentration

in filtered and unfiltered samples is statistically significant.

5.3 Vegetation Studies

Considerable information has been accumulated on the vegetation of the Luquillo Mountains, including the El Verde rain forest (Odum & Pigeon, 1970). Phase I studies are designed to fill gaps in our knowledge of nutrient flows through the ecosystem and to provide a basis for subsequent investigations in phase II.

### 5.3.1 Plot Characterization and 5.3.2 Phenology

Plot characterization and phenology subtasks are being initiated at this time, but preliminary data is not sufficient for analysis. Litterfall and decomposition studies were given priority because of the length of time needed to obtain results and because of their relevance to overall cycling and transport processes.

### 5.3.3 Litterfall

Preliminary results give a mean value for October of 1.73 g/m<sup>2</sup>/day. Plots 1 through 4 varied from 1.76, 1.29, 1.99 and 1.89 g/m<sup>2</sup>/day, respectively.

The leaf fall contributions of the most important species are shown in Table 3. The 1 (highest) through 5 ranking of the species is based on grams of leaves/m<sup>2</sup>/day by plot and overall means.

Considerable time was required to establish the litterfall baskets in the forest. AS a result, only limited data are presently available and no conclusions are possible. The sampling program has been fully implemented and adequate data will be available for analysis in the phase I final report.

239.

---Page Break---

aM loTE

---Page Break---

Plot 1 through 4 contributed 0.04, 0.04, 0.002 and 0.09 g/m<sup>2</sup>, day of flower and 0.07, 0.04, 0.06 and 0.05 g/n<sup>2</sup>, day of fruits respectively for the month of October, 1980,

The species contributing the greatest amount (g dry weight) to the October Flower component of litter were:

Plot 1. *Croton poeeflanthus*

Plot 2, *Honalium racenosum*

Plot 3. *Homalium racenosum*

Plot 4, *Byrsonena coriacea*

#### 5.3.4 Leaf Decomposition

This task will begin in the 1981 dry season (January-April) and be continued into the following wet season. These methods have been designed to avoid possible errors found in other studies. Weigert (1970) collected litter from the forest floor and oven dried the samples before replacing them in the forest. This method was not used because oven drying the samples before and during the experiment probably killed the decomposers and altered the decomposition rates and because samples collected from the forest floor (loose litter) are in all stages of decomposition,

The following modifications are expected to yield more relevant information:

\* placing enough bags in the field so that the collected bags would not need to be replaced after drying

© using freshly fallen leaves to detect rates of decom

Position, caloric content changes, and mineral loss

Decomposition rates for individual species will not be measured

as were done in the studies of Ewel (1976), Meigert and Murphy (1970).

It is believed that leaf composites from each plot will yield the

site and species interactions characteristics of each plot and there-

fore give information of any overall site differences in decomposition

41

---Page Break---

## 5.4 Faunal Studies

Substantial progress has been achieved in defining major reser-

voirs and determining principal pathways of nutrients and energy through

this rain forest ecosystem, A comprehensive view of the food web

organization is emerging which will provide valuable insights into the

Functioning and complexity of tropical forests. Some of the preliminary

results have been unexpected and have caused the redirection of some

tasks. It is tempting to integrate our present information, but premature

analysis might be counterproductive to program goals and will be defer-

red until the final report. Individual task results are presented and

discussed below.

### 5.4.1 Invertebrates

Surveys of invertebrates as described in the methods section (4.2) have been delayed because of staffing difficulties. The position has been filled and studies will begin in February, 1981,

Incidental observations and food analysis suggest that complicated and important interactions occur between larger forms (e.g. spiders and whipless whip scorpions) and small species of vertebrates, mainly frogs and lizards.

#### 5.4.2 Amphibians and Reptiles

Habitat searches were performed to determine which species were present in the study area. No new species or unexpected observations were made during seasonal sampling. A compilation of observations made during phase 1, and data from literature on the El Verde site will be presented in the report which follows dry season sampling. Species observed in the Study Area during phase I studies are listed in Table 4.

Results of the wet season amphibian abundance surveys are presented in Table 5, *Eleutherodactylus coqui* was consistently the most abundant species, although many of the individuals observed were juveniles and subadults. *E. wightmanae* was the next most abundant species and the only

hoe

---Page Break---

Table 4. Amphibian and Reptile

Species Observed in the Study Area During  
Phase I Studies.

Order: Anura

Bufonidae

*Bufo marinus*

Family: Leptodactylidae

Species: *Leptodactylus albilabris*

*Eleutherodactylus coqui*

» *pottoricensis*

*Fichnondi*

*hedricki*

*wightmanae*

*mneidae*



REPTILES,

Order: Squanata

Suborder: Lacertilia

Family: Gekkonidae

Species: Sphaerodactylus macrolepis

?claubers

Family: Anguidae

Species: Diploglossus plees

Family: Auphisbaenidae

Species: Amphisbaena cacca

Family: Tguanidae

?Species: critatellue

Suborder: ophidia

Family: Colubridae

*Alsophis portoricensis*

3.

---Page Break---

2 during 1980 vet

:

Abundance

srved

4

<u

0.3

Bufo marinus

labris

Leptodactylus albi-

90x

gas

¥

3

5

4

?Surveys conducted September 2

fy

Fyeard calling in plot.

---Page Break---

other species observed along transects in 211 four plots.

E, hedricks

was heard calling in all plots, but its densities were such that it was encountered on only two transects. Other species are apparently present at very low densities compared to *E. cogui*. Similar results have been obtained during dry season surveys by Dr. Margaret Stewart during her Stites in the same area, and therefore surveys will not be repeated during the 1981 dry season.

Investigation of food habits of Anolis lizards were begun. The minimum number of lizards were collected for the first seasonal survey, but identification of stomach contents is incomplete. The types and frequency of occurrence of food items from the stomachs which have been examined are presented in Table 6. Conclusions from initial sampling are only tentative, but seem to indicate that the same insect orders (Coleoptera, Lipodoptera, Diptera, and Hymenoptera) are important prey items of both species. *Anolis evermanni* appears to consume more Orthoptera than does *A. gundlachi* which may be a reflection of different habitat use rather than a true food preference.

Following dry season sampling, all stomach contents will be re-examined and identified more specifically (where possible). Data will be correlated with insect/invertebrate sampling to detect possible food preferences and energy/mineral flow pathways.

Relative abundance and minimum population density estimates for the two most common lizard species (*Anolis gundlachi* and *A. evermanni*) are presented in Table 7. Minimum density calculations show higher population levels in plot 4. Inspection of the data show that many of the observations were juvenile lizards on understory vegetation beneath forest canopy which was more open than in the other three plots. *Anolis gundlachi* was found to be the most abundant species observed in all plots, while *A. evermanni* demonstrated considerable variability among plots. This may be a result of the small sample size rather than indicating a real difference,

Two additional lizard species were observed along transects *Anolis*  
*Stratulus* and *A. cuvieri*. Both species are primarily found in the forest  
canopy, and this sampling method consistently underestimates their popul-

85

---Page Break---

Table 6. Frequency of occurrence of food items in *Anolis gundlachi* and *A.*

Frequency (2)

Food Item *Anolis gundlachi*>

EE

Phylum: Aschelminthes 7 8

Class: Nematoda

Phylum: Mollusca 4 -

as

Gastropoda

Phyllue: Arthropoda

Class: Arachnida

Order: Araneae a 6

Class: chilopoda

Scolopendra ap. - 4

Class: Insecta

Orders: Orthoptera 1s 40

Isoptera 7 6

Hemiptera 7 -

Neuroptera - 2

Coleoptera 39 ?a

Lepidoptera 30 20

Diptera 56 76

Hymenoptera 10 8

Chordaca

Reptilia

Order: Squamata, - 4

Plant Material 7 4

Yay adults (15 males, 12 fenates)

705 adults (12 males, 13 feaates)

46

---Page Break---

om ore at 168 we vee ve 3004

st 960 « 106 vo ore

ar ey wt ee 902

o sos act ve ere we 012



yaans uosees 19x Bupanp

UT pue \*saouepunge ot

+o eran

?47

---Page Break---

ation density. Vertical transect studies, not included here, indicate  
?that *Anolis stratulus* may be an abundant species in the rain forest study  
area,

?The minimum population density estimates are considerably below the  
absolute estimates calculated by Turner and Gist (1970) for the same area,  
Relative abundance estimates are also at variance with those presented by  
Well (1978) for similar habitat at similar elevations near El Verde, Pree  
Viminary data from other lizard studies at El Verde suggest that time of  
(ey, weather, and season may all affect the activity and indirectly the  
observability of species, These questions will receive further consider-

ation during dry season sampling.

It is important to note that these data apply only to the lower 4-5 m of the rain forest. Vertical transect studies are being conducted which indicate vertical stratification of lizards, and lizard distribution throughout the 18-22m height of the rain forest. Data from these studies will provide additional data on the relative abundance and distribution of lizards in the rain forest.

Relative abundance and vertical survey for the first wet season have identified *Anolis stratulus*, *A. gundlachi*, and *A. evermanni* as abundant species within the study area. Because of the vertical stratification of these species, and because population density estimates have not been obtained for *A. stratulus* (which may turn out to be the most abundant species),

Studies of population turnover and growth rates have been deferred until data from dry season surveys have been obtained,

### 5.4.3 Birds

Initial field work on the avian section of the CEER-DOE mineral cycling and transport study has produced both useful results and a new perspective on the problems of sampling and the appropriate techniques to overcome these problems. Moreover, recent developments in avian censusing methods require a re-evaluation of proposed sampling plans, with the result that more accurate population estimates can be generated. Common and scientific bird names are listed in Appendix C.

48

---Page Break---

Population density information was estimated from transect data, Appendix D contains data from transect censuses taken during July-November 1980. The increased number of observations in October and November over preceding months is the result of an increase in avian activity and the addition of a second observer, who was unfamiliar with the Emlen transect methods. An intensive training program since that time has reduced the amount of inter-observer variability to acceptable levels,

The outstanding feature of the data is the dominance of all plots by 2 single species, the Guanacaja. The Red-necked Pigeon and the Puerto Rican Tody are the next most abundant species in each site, although their relative position changes in different sites. The proportion contributed by each species to the total number of observations remains relatively constant for most species in all sites during July-November. A notable exception is the Red-necked Pigeon, where the number of observations in

any month 4s determined by the occurrence of fruiting trees around which these birds aggregate. In months where a food tree was in fruit near a transect line, observations of this species would be relatively high,

Table 8 gives the total number of observations of each species in each site during the period July-November 1980. The data are corrected for differences in the number of transects and in the total time spent in each Plot. Most species show very similar values for all four plots. Six species (Ruddy Quail-Dove, Zenaida Dove, Puerto Rican Emerald, Stolid Flycatcher, Stripe-headed Tanager, Puerto Rican Bullfinch) are twice as common in Plot 4 as in any other plot. Plot 4 is more disturbed than the other three plots, but until vegetation analysis is completed, any explanation for the increased abundance of these six species in Plot 4 is only speculative,

Fifteen of the 21 species observed were found in all plots. Those species found in fewer than four plots were either migrants or very rare residents. With increased observation time, species lists for each site are expected to approach complete congruence,

Data from mist-net samples in each plot are shown in Table 9. In almost two weeks of netting, only half the species occurring in the sites

49-

---Page Break---

zo zo

ot aT

8:0 9"0

to to

eer cet

ro 10

e

g

s

?0

0

0

ur

0

aro

yt

6-0 Prezomy weary oreng

20 oyong paezyT weoty o3zeng

sts ?uooSt poxpeu-peut

20 ?nog epreuaz

ot oot £0 2a0g T¥BND Appr

zo | 1 v0 v0 ro ney porFe3-pox,

ao Faney ?~sRay? aR THR Ray ?FaRaT

separa fepxra?/spaya /epatg serra /epasg para

y 2018 © 2014 z 21a T 201g

?sfuyaunos uy auede

NF3 Pav saune> 30 soquns 203 peaseaz09 sxuno> iomeues1 roquonoK-KtAe Bupanp EPIIG Jo  
seyleAtoEgo Tel0L

8 oraey,

50

---Page Break---

Table 9. umber of captures and capture rate for Plot 1-6,

?\_,-

Plot 1 Plot 2 Plot 3 Plot 4 Total

Ruddy Quait-Dove 9 66 3 ?

Puerto Rican Baerald 2 2 2 2 8

Puerto Rican Tody ° 2 7 a 10

Pearly-eyed Thrasher 1 2 2 ° 5

Red-Legged Thrush ° ° ° 1 a

Mack-throated Blue Warbier o ° ° 2 2

Ovenbied ° ° 1 1 2

Bananaguie 3 2 ° 1 ?

Stripe headed Tanager ° 1 ° ° 1

Puerto Rican Tanager © 6 o o &

Total Captures 1s 2 eo 85

Number of ets u wo

Munber of days 3 3 ? 3

Ceptures/net~day 0.45 1030.70 0,37

sie

---Page Break---

were captured, This result 1s consistent with other sites in the wet tropics, where vertical stratification makes some species difficult to catch. The only spectes caught in numbers was the Ruddy Quatl-Dove, a Ground feeder. These results suggest that estimation of population den-

sities with warked birds will require the use of canopy nets to capture some species,

Figure 6 presents data on the height of occurrence of seven species.

Data were collected during transect surveys and are sumed for all four Study plots. Figure 6 conpares heights of species in two different foraging Guilds. Both nectarivores (Puerto Rican Emerald Huemingbird and Banana-Guit) and Targe frugivores (Ruddy Quatl-Dove and Red-necked Pigeon) show Neele overlap in their vertical distribution. The former guild may show ?more overlap in the canopy, but insufficient data are available at this time to test this hypothesis,



Sixteen birds of six species have been collected and their stomach contents preserved. eights of these birds and other individuals captured Guring mist netting for Task 1 are surmarized in Table 10, Insufficient observations are available to begin construction of preliminary food webs; however, Titerature sources have been identified that will contribute much cf the information on diet needed for this task.

Nineteen of 60 birds have been captured and are being processed for orb calorimetry. A detaied protocol for processing bird parts has been developed. Fecal materfal, food itens and stomach contents are being accu lmitated as the other tasks progress. Difficulty in capturing two species (Red-necked Pigeon and Puerto Rican Woodpecker) may result in these species being omitted from the analysis.

#### 5.4.4 Marmals

No mouse-size mammals were trapped in 80 trap nights in the lines of Sherman line traps, and none have been observed in or near the study area sm the recent past. The trapping surveys will be repeated on a larger Scale during dry season surveys.

520

---Page Break---

Table 10. . Mean weights (in grams) of seven species caught in mist

nets.

eee

x x sD.

Ruddy Quail-Dove 38.6 16a

Puerto Rican taerald u 3203

Puerto Rican Tody 2 66 05

Pearly-eyed Thrasher 7 96.7 36

Ree-Logged Thrush 4 80.5 28

Bananaquit 7 mas

Puerto Rican Tanager 4 3307

eee

---Page Break---

---Page Break---

Larver terrestrial mammals had not been surveyed as of November 30, 1960, but the Indian mongoose (*Herpestes auropunctatus*) and black rat (*Rattus rattus*) have been observed within the study area. A semipermanent grids as described in the methods, will be established at plot 3 and

near the sea] the near future. This segment of field surveys has been delayed until field team members were immunized against rabies, 'S a major reservoir for this disease in Puerto Rico (Ti

The Indian mongoose  
Jerkel et al., 1952).

Horizontal nets placed in the rain forest (plots 1 and 2, at the Sonadora suspension bridge, and near the field station) and a vertical net erected at the field station captured three species of bats: *Artibeus lanaiensis*, *Stenoderma rufum*, and *Honophylla rednani*. All three species have previously been reported from the area (Tansitt and Valdivieso, 1970). The first two species are frugivorous and the third is ny

nectarivorous, In

SsctTvorous species have not been observed in the rain forest, although some species (e.g. *Pteronotus fuliginosa*) may forage in openings and above the canopy.

-55=

---Page Break---

## 6.0. PHASE I SCHEDULE

Field sampling began in July, 1980 following the selection of Sample plots. In spite of the late starting date for some tasks, a 1 phase I field work 1s scheduled for completion by August, 1981.

A Final status report which summarizes and integrates material from all tasks 1s scheduled for completion by September 30, 1981, Because of the late starting date of some tasks, submission of this report may be delayed slightly. An important part of the report will be an outline of projected phase II research including continued investigations of key organisms and pathways and an outline of planned experimental studies.

Seminars and presentations based on work accomplished in phase 1 are scheduled to start by May, 1981 following the completion of dry sea-

Some field work. The workshop which was originally planned has been postponed indefinitely.

It is anticipated that several scientific publications, based on individual tasks studies, will result from this phase,

Manuscripts will be prepared simultaneously with the final report.

Papers of a more integrative nature will be written immediately following

Preparation of the final report and will be given priority over routine Project work.

-56=

---Page Break---

## 7.0 ORGANIZATION AND MANAGEMENT

The Terrestrial Ecology Division is a unit of the Center for Energy and Environment Research operated by the University of Puerto Rico for the Department of Energy. Field studies are conducted in the Caribbean National Forest from CEER's El Verde Field Station. Chemical analyses and other laboratory investigations are performed at the Terrestrial Ecology Division's facilities in Rio Piedras.

The principal investigator is responsible for program coordin-

ation but also has specific task responsibilities. Each major task is supervised by a division scientist. Figure presents the overall program organization.

Collective decisions are made regarding the program execution

by all members of the scientific staff including the Terrestrial Ecology

Division head, and Ecology Section Head. Ultimate responsibility within

CER rests with the director of CEER Or, Juan A. Bonnet, Jr. who responds

directly to the President of the University of Puerto Rico, Or. Ismael

Almodovar.

---Page Break---

AMLSINHDTWLNSNNOMIANS

s3ionus

aWwaows

o2902°9

S310NAS NoILYL393A

poem vein

Aina 1p s2uem07 20

95 Meoea]



ae ?avvog

iT waaD

¥ uonn 4)

woiberd Wedsv01, pus Bun19k3 is9104 u

8946 10) sumvons swew

}OUOM 4339 z 91

---Page Break---

## 8.0 LITERATURE CITED

Bannister, B.A, 1970. Ecological life cycle of *Euterpe globosa* Gaertn.

Gh. 88 18 HT. Oda and R.F. Pigeon, eds tropieay rain forest.

U.S. Atonfe Eneray Comission: NTIS, Springfield, Va.

Bray, J.R, and E, Gorman. 1964. Litter production of forests in the

world. Adv. Ecol. Res. 2:'101-157,

Burton, T.M. and 6.E. Likens. 1975. Eneray flow and nutrient cycling

in salamander populations in the Hubbard Brook Experinental Forest,

New Hampshire. Ecology 56: 1068-1080.

Clark, L.R.y Po. Geier, R.D. Highes and R.F, Morris. 1967. The ecology

of insect populations in'theory and practice. Methuen, London.

Coleman, C.O, 1970. Nematodes in the litter and soil of El Verde Rain

Forest. ?Ch. £7 in H.T. Odum and R.F. Pigeon, eds. A Tropical

Rain Forest. U.S. Atomic Energy Comission. "NTIS, Springfield, Va.

De León, D. 1941. Some observations on forest entomology in Puerto Rico. *Caribbean Forester* 2: 160-163.

Drewry, E.G. 1970, A list of insects from El Verde, Puerto Rico. Ch. 10 in H.T. Odum and R.F. Pigeon, eds. *A Tropical Rain Forest*. U.S. Atomic Energy Commission. NTIS, Springfield, Va.

Ehrlich, P.R. and L.E. Gilbert, 1973, The population structure and genetics of a tropical butterfly. *Heliconius ethilla*. *Biotropica*, 2: 69-82.

Enlen, J.T. 1971. Population densities of birds derived from transect counts, *Auk* 88: 332-362,

Estrada, A. 1970. Phenological studies of trees at El Verde. Ch. 10-14 in H.T. Odum and R.F. Pigeon, eds. *A Tropical Rain Forest*. U.S. Atomic Energy Commission. NTIS, Springfield, Va.

Ewel, J.J. 1976, Litter fall and leaf decomposition in a tropical forest succession in eastern Guatemala, *Jour, Ecology* 64; 293-308.

Gilbert, L.J. 1980, Food web organization and conservation of neotrop-

ical diversity, Chapter 2. ?In M.E. Soulé and B.R. Wilcox, eds.  
Conservation Biology. Sinauer Associates, Inc., Sunderland, Mass.

Gilbert, L.£. and M.C. Singer, 1973. Dispersal and gene flow in a  
butterfly species. *Amer. Nat.* 107: 58-72.

Green, R.H. 1979. Sampling design and statistical methods for environ=  
mental biologists. John Wiley and Sons, New York.

59

---Page Break---

Hairston, N.G., F.E. Smith and L.B. Slobodkin. 1960,  
structure, ?population control and competition,  
421-425,

Holmes, R.T. and F.W. Stages. 1975. Bird community dynamics and  
energetics in a northern hardwoods ecosystem. *J. Anim. Ecol.*  
45: 175-200,

Janzen, D.H. 1970. Herbivores and the number of tree species in tro-  
pical forests: *Amer. Nat.* 105: 801-820,

1971. Euglossine bees as long distance pol-

~~ Vinators GF tropical ?plants, Science 171: 205-206,

1973a. Sweep samples of tropical foliage in

"~ SeCtST?description of study? sites, with data on species abundance and size distributions. Ecology 54: 659-687.

1973b. Sweep samples of tropical foliage insects:

" GYFECts GF Seasons, vegetation types, elevation. time of day ecg insularity. Ecology 54: 687-701,

MacArthur, R.H. and A.T. MacArthur. 1974, On the use of mist nets

for population studies of birds, Proc. Nat. Acad, Sci, 71

3230-3233,

Macquire, B-sJr, 1970. Aquatic communities in bromeliad leaf axils

and influence of radiation. Ch.E-6 in H.T.Odum and R.F. Pigeon,

Goss, gp tropical Rain Forest.? U.S. Atomic Energy Commissions

NTIS, Springfield, Va.

McMahan. E.A- 1970. Radiation and the termites at El Verde. CH. E-8

fe,t:T- Odum and R.F. Pigeon, eds. A Tropical Rain Forest. U.S.

Rome Energy Commission, NTIS, Springfield, Virginia,

» and N.F. Sollins. 1970. Diversity of micro-

© Srthreopds GFEer irradiation. ?ch.e-i1 im fits Odum agd RiP. Pegeon,  
eds. U.S. Atomic Energy Commission, NTIS. Springfield, Va,

Nortorell. LF. 1943. Forests and forest entomology. Caribbean Forester  
4: 132-1342

Mattson, J-N- and O.N. Addy. 1975. Phytophagous insects as regulators  
Of forest primary productivity. Science 190: 515-522,

Moll, A-G. 1978. Abundance studies on the Anolis lizards and insect

populations of altitudinally different tropical forest habiters,  
CEER-11.,

Nevlings Lil. Jr. 1971. The ecology of an elfin forest in Puerto Rico.

Je nett yilongring cycle and an interpretation of its seasonaliey,

J. Arn, Arb,"52(4): 585-613,

---Page Break---

Odum, (H-T. and R.F. Pigeon, eds. 1970. A Tropical Rain Forest. A Study

Of irradiation and ecology at ET Verde. U.S. Atomic Energy Commis

sion, NTIS, Springfield, Va.

Overton, M.S. 1971. + Estimating the numbers of animals in wildlife populations, Chap. 21. In R.G. Giles, eds, *Wildlife Management Techniques*. Wildlife Soc., Washington, ?Ds

Ovington, J.D. and J.S. Olson. 1970, Biomass and chemical content of ET Herde lower mountain rainforest plants, Ch-H<2 in HT. Odum and

B.F. Pigeon, eds. *A Tropical Rain Forest*. USAEC, NTIS, Springfield, |  
Ya

Owens, D.F. 1971. *Tropical butterflies: The ecology and behaviour of butterflies in the tropics with special reference to 18 African species*, Clarendon Press, Oxford. 214 pp.

Paine, R.T. 1966. *Methods of vegetation study*. Holt, Rinehart and Winston, Inc., New York.

Sexton, O.J. R. Bauman and E, Ortleb. 1972, Seasonal food habits of *Anolis Linifrons*.? *Ecology* 53: 182-186.

Sturges, F.W., R.T. Holmes, and GE. Likens, 1974, The role of birds in nutrient cycling in a northern hardwood ecosystem. *Ecology* 66: 149  
155.

Tamsitt, J-R, and D. Valdivieso. 1970, Observations on bats and their ectoparasites. Ch. £-9 in H.T. Odum & R.F. Pigeon, eds. A Tropical Rain Forest. TIS, Springfield, Va.

Tierkel, E.S., G. Arbona, A. Rivera and A. de Juan. 1952. Mongoose rabies in Puerto Rico, Publ. Health Rpts, U.S. Public Health Ser. 67 (3): 24-278,

Turner, F.6. and CB. Gist, 1970. Observations of lizards and free frogs in an irradiated Puerto Rican forest. Ch. E-2 in Wet. Odum and Ror Pigeon, eds. A Tropical Rain Forest. NTIS, Springfield, Vay

einbren, M.P. and @.K. Weinbren. 1970. Observations on the mosquito population in the irradiated forest at £1 Verde. Ch-E-I2 in HT. Odum end RuF. Pigeon, eds. A Tropical Rain Forest. NTIS, Springfield. Va.

Wegert, RG. 1970, Effect of ionizing radiation on leaf fall, decomposition, and litter microarthropods of a montane rain forest. Ch. ted in fT, Odum and R.F. Pigeon, eds. A Tropical Rain Forest, NTIS, Springfield, Va.

Wiegert, R.G. and P. Murphy. 1970, Effect of season, species, and location on the disappearance rate of leaf litter in Puerto Rican rain forest, Ch. Hed in H.T. Odum and R.F. Pigeon, eds. A Tropical Rain Forest,



NTIS, Springfield, Va.

---Page Break---

Weir, 2.5. 1969. Importation of nutrients into woodlands by rocks.

Nature 221:

Wolcott, G.N, 1940, An outbreak of the scale insect *Asterolecanium*

*ustulans* on *maga*, *Montzeuna speciosissima*,? Caribbean Forester

2:67.

- 1946. Factors in the natural resistance of

Woods to Termite stack. Caribbean Forester 7: 121-134,

62.

---Page Break---

APPENDICES

63

---Page Break---

Appendix &

## Calorimetry Data on Anolis Lizards

eS

Species and Dry Roof calories per

Sex Weighe Ash ash-free gm.

- gundlachi gt 1.2676 gne -

"oe é 12,6205, 31.3% 4580.3 cal/gn.

? a 1.6814 27.6% 4031.3

"os e 1.0991 16.2% 4284.2

som e 13275 26.8% 4182.0

2 0.6503 29.4%

"8 g 0.4622 34.6%

g 0.7972 15.9%,

so % 0.701 24.9% \*

wom g 0.8246 13.3%

nom # 0.3337 \_\*

A, stratulus 0.3743 \_

8 2.2751 27.9%

£ 6368 12.5%

"os ¢ 1.n175 15.1, 3927.5

"oe é 0.8866 22.3%, -\*

" 7 \$ 0.6302 23.6% 4895.7

uoo® \$ 0.5689 13.38 4093.8

"os é 1.7375 26.5% 4181.8

"oom é 14528 22.0% 4119.9

"8 ¢ 0.7297 22.0% 40,98.8

"om % 0.6293, 16.9%,

\* Pair of values rejected according to criteria described in Methods section.

?\*\*Insufficient sample for ash measurement.

bie

---Page Break---

Appendix

Carton Analysis of Stream Samples Collected from Cara del Indio

(On October 9, 198

10:47 AM, Plow Level: 2,82 USGS Units.

nee

Surface Bottom

carbon Concentration in mg/? Carbon Concentration in mg/1

Organic Inorganic Organic Inorganic

F v F F v F v

3.43 4.50 6,60 497 337 611

2.07 2.56 3.60 4.50 1473 1,73 60 60

4.87 5.97 6.76 7,50 \$27 7.2

2.00 2.97 5,20 7.20 1.33 200 3.25

2.63 2.97 5,60 7,30 4500 57 6.263

eee

Carbon Analysis of Stream Samples Collected from Cera del Indio

on October 30, 1980, 10:45 AM. Flow Level: 4.46 USGS Units.

\_ |

Surface Bottom

Carbon Concentration in mg/l Carbon Concentration in me/l

organic Inorganic Organic Inorganic

F v r v F v Fou

5.24 8.7% 353 362 5.46 8.56 3,57 3,58

6.08 8.66 3.56 3.56 5.25 9.62 3.28 3,46

3.38 8.00 3,58 3.46 5.76 8,503.60 3.69

47 1.19 3.66 5.50 8.18 3,65 3.80

5.08 875385 3,49 5.46 8.35 3,90 3,85

---Page Break---

Appendix 8 (Cont'd)

Carbon Analysis of Stream Sanples Collected from Cara del Indio

?on November 17, 1980, 11:00 AY. Flow Level: 270 USGS Units.

eee

Surface Botton

Carbon Concentration in mg/1 Carbon Concentration in mg/1

Organic Inorganic organic Inorganic

F v F v r v Fr v

0.806 1.142 5,80 5,89 0.573 0.808 5413 5,23,

0.546 0.859 5.70608 0.417 0,670 6.2357

0.457 0.951 5,36 5,47 0.483 04478 5,80 5.62.

0.561 0.717 5,736.05 0.460 0,473 5,64 5,72

0.569 0.616 5.82 3,73, 0.396 0.433 5.92 5,15

ees

Carbon Analysis of Stream: Sanples Collected from Cara del Indio on

Novenber 25, 1980, 10:45 a4. Flow Level!

Surface Botton

Carbon Concentration in ag/1 Carbon Concentration in mg/2

organic Inorganic organic Inorganic

r v F v F v ¥ v

3.51 3m - 335 hae -

3.01 349 - 276 0 2b + -

3.88 23a - 2M 220 -

7.66 Bm : 2.22 hehe -

3.37 23000 = : 8.87 10.62 = -

---Page Break---

Red-tailed Hawk *Buteo jamaicensis*

Ruddy Quail-Dove *Geotrygon sottana*

Zenaida Dove *Zenaida aurita*

Red-necked Pigeon *Columba squamatus*

Puerto Rican Lizard Cuckoo *Saurothera vieilloti*

Puerto Rican Emerald *Geothlypis trichas*

Puerto Rican Tody *Todus mexicanus*

Puerto Rican Woodpecker *Melanerpes portoricensis*

Stolid Flycatcher *Mniotilta stolidus*

Pearly-eared Thrasher *Margarops fuscatus*

Red-legged Thrush *Minocichla plumbea*

Puerto Rican Vireo *Vireo latimeri*

Black-whiskered Vireo *Vireo altiloquus*

Black-anevwhite Warbler *Dendroica caerulescens*

Louisiana Waterthrush *Seiurus notacitla*

Bananaguit *Cooreba flaveola*

Mve-hooded Euphonia *Euphonia musica*

Stripe-headed Tanager *Spindalis zene*

Puerto Rican Tanager *Nesospingue speculiferue*

Puerto Rican Bullfinch *Loxigilla portoricensis*

Latin names according to Bond, Jane:

Collins, London. 256 pp.

1979. Birds of the West Indies.

-67-

---Page Break---

sty oer set 65 6s



si00suea3 uw 09ç 30 zequnu Te201,

suoyaeearssqo jo 9m2 Tel0L

suofaeazosqo Te30,

WoursTIng weory o2z0ng

aefeuey orory oaz0ng

aefeury popeay-ody35

?eruoydeg pepooy-ong

arnbeusueg

STalHH ona porwosy2?>Te

yenay3z0304 wueTETRCT

O8SEA pOIOASTUA?DUTE

aowoa"o4ts PFTOIS

Aoysepooy wPITE 039g

Apo], ww>7y o330ng

Presouy weary o33ang

? ooxong paezyT ueDry o2zaNg

ba «ons ?w038r paxoe0-po¥

@ woe a0g THEN Apna,

8 ney POTFea~pow

apeaar oy nF 30s ay

sstsaqavoed

WF T8309 02 woFanqzaaueD auaDzeg \*T JOG UF yIUON/sezD?ads YORE jo suoTreArDEqo Jo  
eden

@ xipuaady

? ? ϕ

68

---Page Break---

oer oor wt sor ut

m9 Sur eve 68 u se

sao0suesa © oo? 30 49qmnu TeICL

?voraearasqo jo our2 Teac

suoraearasgo [e201

cept DT omy

very (or0s)9e (9"er)LT

cme

(sens

TT

(p,2009) @ sypuaday

WDUTST TNE wore or39N

?aoSeun], uvoyy o3zeng

aoBeury papray-adrzas

ruoudng papooy-2nrE

0Y>I"OATE PITOIS

amypadpooon ueoTy 032903

eH POTFEI~PoH,

seysoqquazed

---Page Break---

oz > > ? » , s2oseuva3 w ggg 30 s9qenv Te20z

ore oT cer 66r ot ast suoyaeaz9sqo 30 942 Tel0]

ws eT sor 03 tw 6 suotaearesgo Te30y

n ame

£

a wne

a

t

esz(e6ND99, (ets) Te (G\*00)9e

6 wae

tr nz ot

t

t

zS woe

3

z

\*

t wor

t

t eH paTTeaepow

aR "ar

yreayauazed

BF sw103 03 woFINGyzaUeD aUB2I" ?ç OTE UF YauOH/soToeds yoee 30 suOTS 30 z9qenk

(P,3009) @ xxpueddy

---Page Break---

© 00? 30 39qunu Tex0x

ey st wer ce est oT ?suoyaeaz9sq0 Jo 8ur3 Tel0,

ots ver, est & woz suoraeezeego [e320]

sre oz Ht CorZDE Hour sTTOe weary o2z9ng

?9 ze Goer (won wot aedvuny uesry o3eng

Uy GE BOT se8eusy, popvay-adr335

90 ? ? wruoydng pepooy-arra

Y'SE TAT (a\*9z)6y (I°ED)99 (B\*¥E)ST (9reRI¥Z (BrESDEL aynbeueury

a91g1ey 92TH Pur 9DeTE

?anyg peavozu2?¥>0Te

?yonaysa93eH eueserNoT

er a emer 025A POlaASTUM OPTS

z0 1 oaz¥a uRoyy 020g

nt ? cer (err ysnauL pes ¥aT-pat

as oe ems "6s sousesul poso-Ktareg

nt ? z0uD3e24Td PETOIS

ez st aeypadpoog weaty 0339ng

s' ? wor ?p01 wary o2zaN

Te oT preaoag weoyy o2zeng

80 Y ory oazeng

zor 66 rene wooStg poDaH poe

90 c ?9aog epreusz

ste ar woz ?40g Tx¥MD Appr

0 z MEH POTEEI=Pou



SOL ae oy

saysoqauozed

UF ste203 02 wor3ngza3¥09 qusDI0q

\*y 20ta ur yauow/sarseds y>ee 30 suoyaeaxseqo Jo 199ENN

(P3409) @ x7puaday

-n-

---Page Break---

---Page Break---