

CEER-T-200

cepr-7-200

Effect of Sunlight and other Factors on the
Population Dynamics of Coliform Bacteria

by

Cindy Ginés sanchez

?Terrestrial Ecology Division,

Center for Energy and Environment. Research

Submitted to the School of Environmental Health,
University of Puerto Rico, Medical Sciences Campus
in partial fulfillment of the requirements for
the Master of Science Degree

+ december 10, 1984

CENTER FOR ENERGY AND ENVIRONMENT RESEARCH

---Page Break---

Effect of Sunlight and Other Factors on the
Population Dynamics of Coliform Bacteria

by

Cindy Ginés sanchez

Terrestrial Ecology Division,

Center for Energy and Environment Research

Submitted to the School of Environmental wealth,

University of Puerto Rico, Medical Sciences Campus

in partial fulfillment of the requirements for

the Master of Science Degree

December 10, 1984

---Page Break---

Bobby, Sandra,

y Kenny

---Page Break---

ACKNOWLEDGEMENTS

So many people to thank! People without whose help and encouragement this work would never have been completed. First I would like to thank the many workers at the Caglias Sewage Treatment Plant, Caglias Police Department, and the sand quarry who helped make the samplings possible. I would also like to thank the personnel at the Terrestrial and Marine Ecology Divisions of the Center for Energy and Environment Research for allowing me to borrow equipment and use their facilities. Special thanks to my good friends in the Center for their help and support, Carlos Figueroa, Carlos Ramos, Lourdes Prieto, Norma Ortega, Roberto Trinidad and Luis Ivin Rosa.

Finally, I would like to thank the many friends who donned boots and gloves to help me with the samplings, among them Oscar Cruz, Lillian Coli, Eliezer Bermidez, and Roberto Ginke.

---Page Break---

TABLE OF CONTENTS

List of Tables. ee ee

List of Figures. 6 6. ee eee ee odd

List of Abbreviations. 2... 1 2 ee iv

Rbstract . ee

Introduction. . ee ee ee D

objectives. © 6. ee 8

Literature Review... . ee ee ee

Materials and Methods... 2... - sod

Results. 6 ee ee ee ee DD

Discussion... 2 6 ee ee ee 8

Conclusions and Recommendations. . . . - . . 35

Literature Cited. 2 6 2. 1 ee ee 2 6

Appendix 2 2 2 ee ee ee

---Page Break---

LIST OF TABLES

Water quality summary for Rfo Cagiiitas

and El Caflo

Coefficients for the correlation of the

physico-chemical parameters with the

logarithm of fecal or total coliforms

ANOVA used to examine the effect of the stream segment, the hour of sampling, and the group of bacteria on the change in the logarithm of coliform bacteria between adjacent stations

Sampling results.

20

22

27

a

---Page Break---

LIST OF FIGURES

Patterns of bacteria mortality

Effect of exposure to sunlight on total

coliforms

Location of the sampling stations

Correlation of chlorine with the logarithm
of coliforms in the sewage treatment plant
outfall

Total coliform density in the Cagilitas
River and El Cano

Fecal coliform density in the Cagilitas
River and El Caflo

Correlation of the logarithm of the total
coliforms with the logarithm of the fecal
coliforms

1s.

23

24

28

---Page Break---

LIST OF ABBREVIATIONS

Abbreviation Meaning

DO Dissolved Oxygen

FC Fecal Coliforms

Log_e Logarithm of the Fecal Coliforms

Log_e Logarithm of the Total Coliforms

TC Total Coliforms

T_w Water Temperature

STP Sewage Treatment Plant

---Page Break---

ABSTRACT

Effect of Sunlight and Other Factors on the Population

Dynamics of Coliform Bacteria: Francisco Folch Castafier, Ph.D., Principal Advisor; William fi. McDowell, Ph.D., Associated Advisor; Iraida Robledo, M.S., Associated Advisor.

The effect of sunlight on the population dynamics of coliform bacteria was studied in a stream receiving the effluent from Cagliari's Sewage Treatment Plant (STP). The effects of nutrients and chlorine were also studied. Five stations were established in El Calflo (into which the STP discharges) and Rio Cagitas (which merges with Rio Camo approximately 0.5 kilometers downstream from the STP discharge). Samples were taken during the daytime and nighttime, and dissolved oxygen, water temperature, light intensity, and chlorine were measured in the field. Samples were taken back to the laboratory for analysis of fecal and total coliforms, nitrates and nitrites, ammonia, and

The results of this study show that the water quality of these streams is extremely poor. Both the fecal coliforms and dissolved oxygen had a high rate of non-compliance with the water quality regulations. Moreover, Rio Cagitas, which does not receive a discharge from a STP, usually had a higher density of coliform bacteria than

EL Cano, whose waters are composed almost solely of the effluent from an STP. A multiple regression-correlation analysis demonstrated that there was no significant effect of either the nutrients or chlorine on the density of either coliform group.

An ANOVA analysis showed that the stream segment significantly affected the change in the density of volifoums between adjacent stations, but the hour of sampling did not. ?This result shows that segments of the streams had different rates of inccese or decrease in the density of bacteria, probably due to the fact that Rio Cagiiitas had a higher density than £1 Cafe. The fact that the hour of sampling did not have a significant effect

sows that there was no difference between daytime and nightie samplings, therefore suggesting that sunlight did not cause substantial coliform die-off in these streams.

---Page Break---

INTRODUCTION

?The use of non-pathogenic bacteria as indicators of pollution with pathogenic bacteria has generated much interest in the population dynamics of coliforms once they are released into streams, lakes or the ocean. Public health officials need to be able to predict the population dynamics of these bacteria in the environment in order to determine what uses can be given to a body of water that receives fecal pollution, In Puerto Rico it is necessary to predict coliform population dynamics because the high human populational density creates a water pollution problem that is increased by industrial waste discharges. The small number of water bodies fit to serve as water supply sources and as recipients of wastes, makes water reuse a necessity that amplifies this problem.

Based on a review of the literature Chamberlin and Mitchell (1978) concluded that sunlight is the most important factor in the mortality of coliform bacteria released into the ocean. Exposure to sunlight can kill over 90% of a population of total coliforms in @ hours (Grigsby and Calkins, 1980). Sunlight therefore appears to be an effective bactericidal agent that is not being utilized optimally to disinfect sewage effluents that are

---Page Break---

released into the environment. However, other factors have also been found to affect coliform survival. The chlorine applied to sewage treatment plant (STP) effluent kills a large proportion of bacteria in the effluent (Camper and MeFeters, 1979), and the density of these bacteria can also be affected by the concentration of nutrients in the water (Hazen and Aranda, 1980).

Therefore, it is obvious that the survival of bacteria released into the environment may be determined by more than one factor. This experiment was conducted to assess the effect of sunlight on the survival of fecal and total coliform bacteria (FC and TC, respectively) released by @ STP into a small, tropical stream in Puerto Rico. Samplings were carried out at day and at night to determine the effect of sunlight on their mortality pattern. The effect of other factors, specifically chlorine and nutrients, was also examined.

---Page Break---

OBJECTIVES

The purpose of this study is to assess the effect of sunlight on the population dynamics of total and fecal coliform bacteria, and to assess the effect of other

factors, specifically chlorine and nutrients.

---Page Break---

LITERATURE REVIEW

Coliform Population Dynamics

Measurement of Coliform Population Dynamics

The magnitude of the die-off of a population of bacteria once it is released into a body of water is measured using Chick's decay coefficient, k (hour), which is a function of the initial density of coliforms (N_0),

the

final density (N), and the time elapsed (t , hour)

(Chamberlin and Mitchell, 1978):

$\log (N_0/N)$

a

t

The final density of bacteria must be corrected for dilution of the plume, and this procedure is explained in John (1978) and Hubbard, et al. (1982).

?This equation is based on the assumption that bacteria released into an unfavorable environment will experience first-order die-off. This die-off pattern is characterized by the death of an equal proportion of the remaining Population of bacteria in a given time period (See Fig. D. Woweever, bacteria populations do not always follow this

pattern,

---Page Break---

Population Dynamics in Temperate Regions

and lakes

Studies carried out in temperate stre show a great variability in k values. In their review of the Literature Chamberlin and Mitchell (1978) report that k

values range from 0.010 to 1.0 and generally fall between 0.015 and 0.20 hour⁻¹. Another author (Velz, 1970) reports that during the warm weather period k values for temperate rivers ranged from 0.46 to 0.96. The discrepancy between these authors is not easily explained. One possible reason is that Chamberlin and Mitchell (1978) corrected the k coefficient for the dilution of the effluent, but Velz (1970) apparently did not. The most likely explanation is that the studies they cited lasted different time periods.

The studies used by Velz (1970) lasted 1 to 2 days and were

x

s

"wo

i

1» ? _

todo at Yodo

Doys

Fig. 1. Patterns of bacteria mortality. 1st indicates

first-order and 31a indicates third-order die-off. Adapted from Mancéni (1978).

---Page Break---

carried out during the summer, but Chunberlin and mitenell (1978) do not state how long the studies they reviewed lasted or in what season they were done.

Based on a review of the literature and on his own sampling, Velz (197C) concluded that the most common

pattern of coliform mortality is third-order die-off. This

pattern consists of rapid initial die-off followed by much slower mortality (See Fig. 1), He found that 99.518 of a coliform population had a half-life of 0.64 days, and the

other 0.49% had a half-life of 0

5 days. Therefore, he

concluded that the mortality pattern was basically first-

order since such a large fraction of a population of

bacteria die during the first day in a river.

Population Dynamics in Puerto Rico

The dynamics of coliforms appear to be different in the tropics than in temperate regions. k values are much

higher, and regrowth of bacteria has been reported, one

study of coliform mortality carried out in Puerto Rico

found k values that ranged from 0.8 to 2.9 day⁻¹ (19.2 to 23.2 hour⁻¹) (Munoz, et al., 1969). The k values reported

were much higher than those reported for temperate regions

and demonstrate that die-off in the tropics can be much

quicker than in temperate regions. However, the same study

also reports that in two Puerto Rican streams coliform

density increased instead of decreased. Niambn and Hazen

(1983) also report that coliforms in an effluent plume

---Page Break---

dumped into the ocean showed an initial increase followed by a decrease in density. ?These findings may be explained in part by the fact that in both cases of regrowth the polluting material was the organic-rich waste material from sugar cane. This effluent may provide nutrients necessary for bacteria reproduction that are not provided by sewage effluent. Another possible explanation is that in the tropics phytoplankton is involved in the control of population levels of coliforms (Hazen and Aranda, 1980).

Population Dynamics of Pathogenic Bacteria

It is important to remember that the population dynamics of pathogenic bacteria do not necessarily have to be the same as the dynamics of coliforms. Several studies

have demonstrated that pathogens have greater resistance to lethal agents than coliforms (Chamberlin and Mitchell, 1978; Jagger, 1981). In their study of bacteria in an

industrial effluent, Biamin and Hazen (1983) found that the density of FC was not correlated with the density of *Aeromonas hydrophila*, a pathogenic bacteria, Therefore, care must be taken when making generalizations of bacteria mortality based on studies of coliforms.

---Page Break---

Factors That Affect Coliform Population Dynamics

Sunlight

Based on an extensive review of the literature Chamberlin and Mitchell (1978) concluded that sunlight is the most important factor in the mortality of indicator bacteria released into the ocean, They found a significant linear relationship between the decay rate of coliforms and the depth-corrected intensity of sunlight. Evidence of sunlight's lethal action to bacteria in freshwater was provided by Grigsby and Calkins (1980) who placed samples of effluent from a sewage treatment plant in tanks exposed to the sun. They found that the density of TC decreased steadily during the day and increased slightly at night (See Fig. 2). Several other in vitro experiments have

Time (hours)

Fig. 2, Effect of exposure to sunlight on total coliforms.

The bar represents daytime (empty) and nighttime (filled) exposure. Adapted from Grigsby and Calkins (1980).

---Page Break---

onstrated that sunlight causes a decrease in the density of coliforms (Chojnowski, et al-, 1979; Fujioka and Narikawa, 1982; Tuveson and March, 1980).

Therefore, there is widespread evidence that sunlight causes mortality in vitro, but there have been no experiments to our knowledge to test the effect of sunlight on coliform mortality in sewage plumes, Proving this effect may be difficult because the turbulence and turbidity of plumes may attenuate sunlight's lethal action. However, at least indirect evidence has been presented by Fattal, et al. (2983) who detected a negative correlation between the mean monthly FC density in a sewage plume and the mean

monthly duration of sunlight.

The ultraviolet (UV) component of sunlight is at least partially responsible for its lethal effect. All three components of UV light {near (λ= 320-400 nm), mid (λ= 290-320 nm), and far (λ= 200-290 nm) UV} can kill bacteria (Peak, 1983). Their modes of action are different. The mid UV component causes mixed results, but the far and near UV cause the same end result:

death of the bacteria due to DNA damage (Peak, 1983). The near UV can also cause sub-lethal damage to the permeability of the cell membrane and protein synthesis apparatus (Jagger, 1961).

Chlorination

Chlorine is the most commonly used disinfectant of waste water effluents, It rapidly reduces coliform

---Page Break---

Densities by as much as 90% (Camper and McFeters, 1979).

However, recent experiments have shown that this decline is only temporary and, that after a lag period, the density of coliforms begins to increase. Following exposure to chlorine the density of indicator bacteria decreases for an hour (Camper and McFeters, 1979), but after this initial decline the density can increase for as long as five days (Kinney, et al. » 1978).

The explanation of this "regrowth" lies in chlorine's mode of action. A large proportion of the bacteria exposed to chlorine are not killed; they are injured sub-lethally. Although chlorine damages the cell's genetic material and protein synthesis apparatus (Haas and Engelbrecht, 1980), the greatest damage occurs in the cell membrane. Pores are opened in the membrane that disrupt active transport and allow low molecular weight material to escape the cell (Haas and Engelbrecht, 1980). The net result is that the bacteria is not dead but is injured to such an extent that it is unable to grow on selective media such as m-FC and m-endo. Proof of this hypothesis was provided by an experiment in which the absorbance of a suspension of chlorine-injured coliforms was measured simultaneously with plate counts. The former remained unchanged while the latter decreased (Camper and McFeters,

1979). Given time, the cells repair the damage and are

again able to grow on the plating media, creating the

illusion of regrowth?

10

---Page Break---

ALL research involving indicator bacteria must take into consideration the fact that the most commonly used plating media do not detect these sub-lethally injured cells. m-endo can recover 95% of the uninjured and 668 of the chlorine-injured cells, and n-PC can recover 1058 and

7%, respectively (McFeters, et al., 1982).

Nutrients

Little emphasis has been placed on the role of nutrients on the population dynamics of coliforms in the environment. In their review of factors influencing coliform mortality, Chamberlin and Mitchell (1978) concluded that nutrients could explain only a small fraction of coliform die-off. However, several experiments

have demonstrated that a correlation can exist between nutrients and coliform density. Research conducted in a Swedish estuary showed that the density of FC was positively correlated with total nitrogen and nitrate (Hirn, et als, 1980). Razen and Aranda (1980) found that the density of FC in several stations along a Puerto Rican stream was positively correlated with ortho-phosphate and nitrate concentrations. A study of the population dynamics of bacteria in a plume of organic-rich industrial effluent dumped into the sea showed that the density of the bacteria *Acreowsnes hydrophila* was positively correlated with total phosphorous, total organic carbon, and orthophosphate;

whereas the density of FC correlated only with conductivity

n

---Page Break---

(Biam?n and Hazen, 1983). Since a relationship between

bacteria density and nutrients does not always exist, and since the nutrients involved vary, it is possible that the

effect of nutrients is indirect or only one of many. One possibility is that phytoplankton may be modifying the

effect of nutrients on coliforms (Hazen and Aranda, 1980).

other

Various other factors have been proposed to cause coliform mortality, including predation. In one experiment, predation was found to remove 10^7 coliforms in 10 days (McCambridge and McMeekin, 1979), but in another experiment predation did not cause significant die-off (DeLeval and Remacle, 1979). This incongruity is probably due to the different methodologies used. The latter experiment removed the protozoans by filtering the water through a 0.8 μ m membrane, but the former experiment demonstrated that it is this component of the fauna that is most active in predation,

Sedimentation could also affect coliform population dynamics. *E. coli* and *Vibrio cholera* can survive for considerable periods of time and even reproduce in sediments (Hood and Ness, 1982). Furthermore, activities such as dredging the bottom of a stream can resuspend coliforms bound to the sediment and increase their density in the water column (Grimes, 1980). Therefore, sediments

jay act as a sink or a6 a source of coliforms.

12

---Page Break---

Still other factors have been proposed, including heavy metals, bacteriophages, algae, and bacterial toxins. However, their effect on the population dynamics of bacteria is deemed to be minimal (Chamberlin and Mitchell, 1978).

a3

---Page Break---

MATERIALS AND METHODS

Location

The source of fecal bacteria for this study was the city of Caguias' STP, The trickling filter plant received mostly domestic wastes at an average flow of 6.9 Mod (PRASA, 1984). The plant discharged the treated effluent into a channel locally called El Cafio (See Fig. 3). During stable flow conditions the plant's effluent makes up the entire flow of El Cafio, which runs through 0.5 km of

Pasture before merging with Rio Caglitas. Downstream from the point of merger with El Cafic, Rio Caglitas flows by a sand and gravel quarry; however, this activity does not seem to affect the river's flow significantly.

Both Rio Caglitas upstream from El Cafic, and the stP's effluent have highly variable water quality. At times both the river and the plant's effluent bear almost raw sewage and at other times the water they bear are moderately Clean. For other studies of this river see Quiffones-Marquez (1980) and Font, et al. (1972).

Sampling Procedure

Five stations were established in a 1-km stretch of the Caguitas and El Cafic (See Fig. 3). Station 1 was

4

---Page Break---

?resiocntia seca ©

STREET wemeer

ster ©). sawpune stariow

EL eNE rc war MARKER

Fig. 3. Location of the sapling stations. Cagilas

Sewage Treatecnt Plant (SIP in the eup) ds located in

eastern edge of the city. Adapted from USGS

---Page Break---

located at the point of discharge of the STP into El Caflo.

Station 3 was located in Rio Caglitae upstream from its

merger with El Cafio and is equivalent to the Geological

Survey's water quality station number 50055250 (Curtis, et al., 1984).

Samples were taken once a week beginning in March

1984, Five daytime samplings were followed by three

nighttime samplings. All samplings were carried out in stable flow conditions beginning at 9 a.m, and 9 p.m, respectively. The order in which the stations were sampled varied but was generally: 3, 1, 2, 4, 5.

Two samples were taken at each station, A sample for microbiological analysis was taken in a sterile bottle with 1 ml of 10% sodium thiosulfate to neutralize residual chlorine (APHA, 1980). A sample for chemical analysis was taken in a plastic bottle. The samples were taken simultaneously 10 cm from the water surface and were Protected from sunlight immediately. All samples were

Placed on ice as soon as possible (but never more than five minutes after sampling) and were kept on ice until they were analyzed,

Several other parameters were measured in each station at a depth of 10 em. Dissolved oxygen (OO) and water temperature (Temp) were measured using a Model 57 Meter (Yellow Springs Inst. Co.). Light intensity was measured using an underwater photometer (Protomatic). Total Chlorine was measured in Station 1 with a Model Ch-66 meter (Hach Coo.

16

---Page Break---

Coliform Enuoeration

Coliform densities in the eamples taken for microbiological analysis were determined using the membrane filtration method as described in Standard Methods (APHA, 1980). M-endo and M-Fc broths (Difco) were used to determine total coliform and fecal coliform densities, Tespectively. These media were filter cterilized through 0.20 4m pore diameter membranes (Type SM, Sartorius Co.) and dispensed into sterile petri dishes with absorbent pads (Millipore Co.)

A sample was chosen randomly for analysis. Using sterile phosphate buffer (APHA, 1980), two dilution series (one for each medium) were made. The final filtration volume was always 10 mL, and these dilutions were filtered in less than 30 minutes. A sterile filtration unit and 0.45 µm pore diameter membranes (Type HA, Millipore Co.) were used. The four dilutions for one medium were filtered beginning with the most dilute. The filter was then sanitized by rinsing it with ethanol, was allowed to air dry, and was rinsed with sterile buffer before beginning the remaining filtrations for that station. The same procedure was followed for each station, and the maximum time elapsed between collection and examination of the samples was 8 hours.

TC were incubated at 35°C for 24 hours, and red colonies with green metallic sheen were counted. FC were

incubated at 44.5°C for 24 hours, and blue colonies were counted,

Chemical Analysis

Samples were analyzed for nitrite and nitrate, ammonia, and chloride, using automated analysis (autoAnalyzer II, Technicon Instrument Co.) as described in Standard Methods (APHA, 1980). The samples were prepared by filtering them through Whatman filter paper (W & R Balston Co.) and then through 0.45 μm pore diameter membranes (Type BA, Millipore Co.). The filtered samples were diluted 1 in 10 using distilled-deionized water (Wanopure, Barnstead Co.) before analysis.

A portion of each diluted sample was stored at 4°C for determination of nitrogen as nitrate and nitrite ($\text{NO}_2^-/\text{NO}_3^-$) and, chloride ion (Cl^-) (holding time < 5 days, and 3 weeks, respectively). A second portion was acidified by adding concentrated sulfuric acid and stored at 4°C for analysis of nitrogen as ammonia (NH_3) (holding time < 6 days).

Data Analysis

Data analyses were done using the SPSS statistical

Package (Nie, et al., 1975).

18

---Page Break---

esuuts?

Water Quality

A summary of the values obtained for several indicators of the water quality of Rio Cagilitas and £1 Cao appears in Table 1. As expected, the high densities of FC and TC, low concentrations of DO, and extremely high concentrations of Nig*, corroborated that both streams receive sewage effluent.

?Two of these parameters (FC and DO) had a high rate of non-compliance with the applicable water quality regulations (EQB, 1983). Whereas the standard is that no more than 20% of all samples taken in a given location shall have more than 40 CFU/mL of fecal coliforms, 80% of the samples in this study exceeded that value. Over 908 of all samples analyzed for DO had a concentration lower than that stipulated by the standard, 4.0 ppm. on the other hand, both C1~ and No2"/NO3 had 100% compliance.

1. All volves obtained in this study appear in Table 4 in the Appendix.

a9

---Page Break---

Table 1, Water quality summary for Rio Caglitas and El Cafic. Results based on all samples taken during this study.

OO

re ena) yes swoo | aan aes

Nag* (ppm) | 9.9 = 14800 } 1350 £ 2971

1, Measured only in the STP outfall (station 1).

20

---Page Break---

Effect of Physico-Chemical Factors

A multiple regression-correlation analysis was carried

out to detect any correlations between the logarithm of either coliform group and the physico-chemical factors studied. The correlation coefficients appear in Table 2, but there was no significant multiple correlation with the logarithm of the bacteria ($p=0.05$). In addition, the

logarithm of neither coliform group in the STP w

significantly correlated with the concentration of chlorine

in the STP (See Fig. 4).

Coliform Population Dynamics

Although the number of bacteria fluctuated greatly with each sampling, a prevalent populational trend was evident (See Fig. 5 and 6). In general, there was a decrease in density between the STP and Station 2, which

followed by an increase between Stations 2 and 3 and a Moderate decrease between Stations 4 and 5. However, instead of the expected decrease in density between the first (Station 1) and Last stations (Station 5), there was @ slight increase in density. This increase may be attributed to the fact that Rie Cagiitas (station 3) usually had a much higher density of coliforms than £1 Caflo and therefore greatly increased their density in Stations 4 and 5 (See Fig. 5 and 6).

Several other trends could be detected. A foreseeable

result is that the total coliforms had a higher density of

---Page Break---

Table 2. Coefficients for the correlation of the physico-chemical parameters with the logarithm of fecal or total coliforms. & stepwise multiple regression-correlation analysis with forward inclusion proved that none of the Parameters were significantly correlated to either coliform group ($p=0.05$).

eee

T

1 Dependent Variable

1

r T

1 Logarithm 1 Logarithm

Independent Variable | Fecal Colif. | Total Colif.

ee ee

r r

Dissolved Oxygen 1 -0.60 | 0.48

1 !

?Temperature 1 0.83 ' 0.37

1 '

Light I -0.03 1 0.12

i 1

8027/03? 1 0.44 1 0.64

' 1

way | -0.50 1 -0.68

I '

Discharge 1 -0.34 ' 0.55

' 1

Chlorine 1 ~0.62 1 -0.50

ne as 1

22

---Page Break---

ab wo a a a

Chlorine (ppm)

L

o

®

2

F

© 2

- S lo

aod ab af lod

Chlorine (ppm)

Fig. 4. Correlation of chlorine with the logarithm of coliform:

in the Sewage Treatment Plant outfall. Neither correlation was

significant (p > 0.05)

---Page Break---

100,000,

10,000

t

- 2 1,000 +

c

1

2 100

- ϕ

F

u

/

7

tot,

r > 5 4 3

STAI LON

- Figure 5. Total coliform deasity in the Cogiitas River and EL

Cabs. Vatures re the average Lor each sampling tat ion

SIF one standard deviation for daytime (@) and night ine (@)

---Page Break---

100,000

10,000

100

1 23% 5

STATION

Figure 6. Fecal coliform density in the Cagtlitas River and #1

Cyl. Vutues represent the average for each sampling station

Sith one standard deviation for daytime (e) and might ine (@)

ings

2s

---Page Break---

bacteria than the fecal coliform:

contrary to what was expected, the average density of both groups of bacteria was greater during the daytime samplings than during the nighttime samplings.

An ANOVA analysis was carried out to determine if the stream segment, the group of bacteria, or the hour of sampling affected the change in the logarithm of coliform bacteria between adjacent stations (See Table 3). of the Patterns that were apparent in the graphs only the Gifference between stations was statistically significant

($p < 0.05$), thereby proving that the increases or decreases

in the density of coliforms were not the same for all reaches of the river. Since there was no effect of the hour of sampling, the apparent differences between daytime and Aightime samplings that could be observed in Fig. 5 and 6 Were not statistically significant, Likewise, the apparent @itferences in the change in the number of bacteria measured by the two coliform groups were not substantial. AS expected, the logarithms of the two groups of

bacteria (PC and TC) were highly correlated (See Fig. 7).

The logarithm of TC accounted for 64.5% of the variation of

the logarithm of Fc.

26

---Page Break---

Table 3. ANOVA used to examine the effect of the stream Segment, the hour of sampling, and the group of bacteria on the change in the logarithm of coliform bacteria between adjacent stations. The stream segments were classified into the areas between Station 1 and 2, 2 and 3, 3 and 4, and 4 and 5. The hour of sampling was classified into daytime and nighttime, and the group of bacteria was classified into total and fecal coliforms. The dependent variable was the difference in the logarithm of coliforms between these stations.

Source of sum of Mean Signif.

Variation Squares DP square oF of F

Bour 0.69 1 0.69 0.63 0.432

Segment 29.00 3 © 9167 8.78 0.000

Group 012-2 0112 or1 0746,

?Two-way Interactions:

Segment and Hour 3.19 3 1.06 0.97 0.416

Segment and Group «11743-0158 05302665

Hour and Group 0:78 =i) 0:78 = 02710403,

?Three-Way Interaction:

Segment, Hour and

Group? or Error 0.10 3 0.03 0.03 0.992

Residual 52-83 48 © 1.10

Total 68.45 63 1.40

27

---Page Break---

*(10"0 & 59

TH1e9 waz aya jo wystaeoy 942 yin suse)

or

Se

oe

9°0 * 24) awessyutye Kua

F109 16: 35

83057109 TH294 Sor

SZ OZ

74103

yp sanded

28

---Page Break---

DISCUSSION

water Quality

The results of this study agree with the findings of another study carried out in Rio Cagiitas (Quinones-Miequez, 1980). The values of dissolved oxygen, chloride, end total and fecal coliforms detected in both studies were in the same range. But the values obtained for $\text{NO}_2^-/103^-$ and especially for NH_4^+ were much higher than the values reported by Quiffiones-Marquez. The averages obtained in

this study were 0.41 and 1350 ppm, respectively, whereas the averages obtained by Quinones-mMarquez were approximately 2.2 and 4 ppm, respectively. This discrepancy could be due to the fact that they only sampled the Cagiiitas upstream from its merger with El Cafe (equivalent to station 3 in this study). This station generally had a lower concentration of Niig* than the other stations. Another possible explanation might be found in the large fluctuation in these parameters with each sampling (See Table 4). Whereas in one sampling the NH₄* concentration was 87 ppm, the next sampling it was 2290 ppm. With such large fluctuations, the discrepancy could

29

---Page Break---

be explained. Of course, the best explanation may be that the quality of the river has worsened since Quinones-Marquez studied it.

The high concentrations of nutrients, low

concentrations of dissolved oxygen, and the high densities

of coliforms found in Rio Caglitas and E Caflo corroborate Quifiones-marquez's (1980, p.51) conclusion that "... the quality of its water ranks among the worst in Puerto Rico." The similarity in the values obtained in both studies Proves that the quality of the river has not improved in four years despite a doubling of the s7P's capacity in 1963. In fact, it appears to have worsened as far as Niig* and NO₂/NO₃ are concerned. & disturbing conclusion that can be drawn from these data is the fact that Rlo Cagitas, which Goes not receive STP effluent, is equally as polluted as El Cafio, whose waters are composed almost solely by the effluent from a sewage treatment plant.

Coliform Population Pynanics

The two groups of coliforms, the fecal and total coliforms, behaved as expected in detecting fecal pollution (APIA, 1960). A lower number of bacteria grew on the fecal coliform medium than in the total coliform because it is a more selective medium, On the other hand, the number of bacteria detected by the two plating media were always te because they both measure fecal pollution,

Proportion

x0

---Page Break---

Although there was a significant effect of the stream segment on the density of coliforms detected, it was impossible to detect any clear-cut die-off or regrowth of the bacteria. The decrease in the density of coliforms between Stations 1 and 2 suggests that the bacteria released by the STP experienced die-off. However, the overall trend in these streams was an increase in the density of bacteria, as indicated by the increase between the STP and Station 5. An ANOVA showed that the stream Segment significantly affected the change in coliform bacteria (See Table 3). This result is probably due to the fact that Rio Caguitas had such a higher density of bacteria than El Cafio that it resulted in a change in density between Stations 2 and 3 that was significantly different from the other segments. Therefore, it is impossible to tell whether the increase between Stations 1 and 5 was due to the addition of bacteria by the Caguitas or the regrowth of chlorine-injured cells.

To determine if the bacteria released by the step experienced die-off or regrowth, a mathematical calculation must be carried out to correct for the number of bacteria added by Rio Cagliitas (Velz, 1970). This study attempted to carry out this correction using the chloride that is naturally found in these waters as a conservative tracer, but the calculation was not successful apparently due to the addition of sub-surface flow that diluted the chloride.

a

---Page Break---

Effect of Nutrients and Chlorine

Neither NH_4^+ , nor $\text{NO}_2^-/\text{NO}_3^-$, nor chlorine affected the density of the coliform bacteria. This result does not agree with a study conducted in Puerto Rico that found a correlation between $\text{NO}_2^-/\text{NO}_3^-$ and the fecal coliforms (Hazen and Aranda, 1980). However, the authors hypothesized that this relationship is indirect and is due to a direct symbionism between algae and the coliforms. Their hypothesis is corroborated by other studies (Cole, 1982).

Hazen and Aranda studied a clear mountain stream where there should be an abundant population of algae. Rio Cagiitas and £1 Cafio, on the other hand, which carry a heavy load of suspended and dissolved solids (Curtis, et al.) 1984) which filters out sunlight and may inhibit algae from growing. Therefore, the correlation of nutrients with coliform density should not be expected in these streams.

A surprising result is the lack of an effect of chlorine on the density of the coliforms. It was not correlated with the density of either group of bacteria in the STP outfall. The lack of correlation shows that the Plant's operators applied excessive quantities of chlorine to the effluent. The concentration of total chlorine in the outfall surpassed 3.0 ppm more than once, This excess did not cause a proportionate decrease in the density of coliforms but instead may have caused damage to the stream's biota (Esvelt, et al., 1973).

32

---Page Break---

Effect of Sunlight

Sunlight did not affect the dynamics of the coliforms

in this stream, The non-significant effect of the hour of sampling on the change in the density of the bacteria between stations indicates that there was no difference between day and night samplings (see Table 3). In addition, the average density of both groups was higher during the daytime than during the nighttime (See Fig. 5 and 6). If sunlight had acted as a bactericidal agent as suggested by Grigsby and Calkins (1980), then the coliform density should be lower during the daytime. Therefore, sunlight was not an effective bactericidal agent in this stream,

Sunlight's bactericidal action has only been proven in highly controlled experiments so it is not too surprising that no effect was found in this experiment. The turbidity and turbulence of the stream's water may attenuate the effect of sunlight by filtering out the uv rays and moving the bacteria to the bottom of the stream where the Uv radiation does not penetrate, Tuveson and Hareh (1980) found that clear water filtered out UV rays and attenuated its lethal effect. The reduction of UV lethality should be even greater in these streams that carry a heavy load of dissolved and suspended solids (Curtis, et al., 1984).

This finding demonstrates that the hypothesis put forth by Chamberlin and Mitchell (1978) that sunlight is

the most important factor in the mortality of coliform

38

---Page Break---

bacteria released into the ocean cannot be extrapolated to all streams, El Cafio and Ric Cagiiitas do not behave as expected probably due to the high degree of turbidity of their waters.

34

---Page Break---

CONCLUSIONS AND RECOMMENDATIONS

The results of this study show that there was no significant effect of sunlight on the population dynamics of coliform bacteria released into El Caflo. There were also no effect of nutrients or of chlorine. In addition, the water quality of these two streams was extremely poor.

It would be useless to suggest ways to enhance the bactericidal effect of sunlight when the water quality of both streams is so poor. Sunlight-induced mortality should

be used as a second line of defense against fecal
Pollution; the first line of defense must be the efficient
functioning of the sanitary system, Not only is the sewage
treatment plant not functioning correctly, but the Cagiiitas
Feceives pollution from cources other than the STP that
Give it a higher degree of fecal pollution than £1 Cafe.

The first step in correcting the gross pollution of
thie area is to identity and eliminate the source or

sources of fecal pollution in Kio Cagiiitas. The next step

must be to increase the efficiency of Cagiiias? stP, which
now barely provides primary treatment, Only when this
Plant removes the high load of solids in its effluent can
suntight begin to be a contributing factor in the control
of indicator and pathogenic bacteria.

25

---Page Break---

LITERATURE CITED

APHA. 1980. Standard methods for the examination of water and wastewater, 15th ed. American Public Health Association, Washington, D.C. 1134 pp.

+ Biamén, E.J. and F.C. Hazen, 1983, Survival and

distribution of *Aeromonas hydrophila* in near-shore coastal waters of Puerto Rico receiving rum distillery effluent. *Water Res*, 17:319-326.

Camper, A.K. and G.A. McPeters, 1979. Chlorine injury and the enumeration of waterborne coliform bacteria. *3rd Appl. Environ. Microbiol.* 37(3):633-641,

Chamberlin, C.E. and R. Mitchell. 1978, A decay mode) for enteric bacteria in natural waters. p. 325-347 in

R. Mitchell, ed. *Water Pollution Microbiology*. Wiley and Sons, W.¥.

Chojnowski, K.J., J.L. Mancini, and J.S. Jeris. 1979. Influence of light on coliform mortality rates in fresh water and sea water. *J. Water Poll. Cont. Ped.*

51(10) 2538-2539,

Cole, J.J. 1982. Interactions between bacteria and algae in aquatic ecosystems, *Ann. Rev. Ecol. Syst.* 13:291-314.

Curtis, R.E., S. Guemin-rios, and P.L. piazz. 1984. Water resources data for Puerto Rico and the is. Virgin Islands, 1983. uscS Water Data Report PR-83-1- available from: USGS, GPO Box 4424, San Juany PR 00936.

DeLeval, J. and J. Romacle. 1979. The estimation of bacterial predation by aquatic microfauna, *Water Res.* 13:513-517.

+ 80B. 1983. Puerto Rico water quality standards

regulations. Environmental Quality board Resolution No,
R-83-5-2, ?as amended February 28, 1983, Available fro
OR, ox'11488, Santurce, Px, 00910,

-Esvelt, GA, W.J. Kaufman, and R.B, Selleck. 1973.

Toxicity

Je Watee P

scasuent of treated municipal wastewaters.

V1. Cont, Ped, 45(7) 21558-1972,

36

---Page Break---

a.

12.

13.

aa

as.

16.

a.

19.

20.

a.

Fattal, B., RJ. Vasl, E. Katzenelson, and B.1. shuval.

21983, Survival of bacterial indicator organisms and

gnteric viruses in the Mediterranean coastal waters off

Tel-Aviv. Water Res. 17(4):397-402.

Fon

feghGrr Sh al. 1972. Report of the sanitary
gonditions of the Loiza river and its
Unpublished report. ?Available from: Puerto iyas
Aqueduct and Sewer Authority, San Juan, PR. 9 pp,

Fujioka, R.S. and O.7. Narikawa, 1982. Effect of
sunlight on enumeration of indicator bacteria under
field conditions, J. Appl. Environ. Microbiol.
44(2) 2395-401,

Grigsby, P. and J. Calkins, 1980. Inactivation of a
natural population of coliform bacteria by sunlight.
Photochem. Photobiol. 31(3):291-294.

Grimes, D.9. 1980, Bacteriological water quality
effects of hydraulically dredging contaminated upper
Mississippi river bottom sediment. J. Apple Environ,
Microbiol. 39(4): 782-789,

Haas, C.N. and R.S. Engelbrecht. 1980. Physiological
alterations of vegetative microorganisms resulting from
ghlorination, J. Water Poll. Cont. Ped. S2(7)s1976~
1987.

Razen, TC. and C.F. Acanda. 1981. The relationship between the distribution and abundance of bacteria and water quality in the Rio Mameyes watershed. p. 87-111 in Séptimo Simposio de Recursos Naturales. Available from: Department of Natural Resources, San Juan, PR,

Hirny J... Viljamaa, and M. Raevuord., 1980, The effect of physicochemical, phytoplankton and seasonal factors on faecal indicator bacteria in northern brackish water. *Water Res*, 14(3):279-265-

Hood, M.A. and G.E. Ness. 1982. Survival of *Vibrio cholerae* and *Escherichia coli* in estuarine water and Sediments. *J. Appl. Environ. Microbiol.* 43(4):578-564,

bubbard, f., FA. Kilpatrick, L.A, Martens, and J.P.

Gilgen. Je. 1982. Measurement of time of travel and

Sispersion in streans by dye tracing. U.S. Geological
Survey Techniques of Water Recources Investigation, ch.
AG, Mook 4. Available from: Branch Distribution, Wscs
1200 South Fads St., Arlington, VA

Jagyer, 1981. wear-UV radiation effects on
microorganisms. Photochem, Photbiol. 34:761-768,

w

---Page Break---

22.

24.

25.

26.

27.

28.

29,

30.

a.

32.

33

John, P.H. 1978, Discharge measurement in lower order streams, *Int. Rev. ges. Hydrobiol*, 63(6):731-755,

Kinney, E.C., D.W. Drummond, and N.B. Hanes. 1978. Effects of chlorination on differentiated colitons groups. *J, Water Poll. Cont, Fed*, §0(10):2307-2312,

Mancini, J.L. 1978, Numerical estimates of coliform Rortality rates under various conditions. *J. Water Poll. Cont, Ped*. 50(11):2477-2483.

McCambridge, J. and T.A. McMeekin. 1979. Protozoan Predation of *Escherichia coli* in estuarine water

Water Res. 13:659-663,

McFeters, G.h., S.C. Cameron, and M.W. LeChevallier.

1982. Influence of diluents, media, and membrane filters on detection of injured waterborne coliform bacteria, J. Appl. Environ. Microbiol. 43(1):99-103,

Muoz, Roy M. Martinez, A, Rodriguez, M. Pedraja, R.

Seda, and C.m. Chan. 1969. The self-puri-

ties of polluted streams in Puerto Rico. Final Report,

WRRRI Project No. A-006-PR. Available from: Water

Resources Research Institute, Univ. Puerto Rico,

Mayaguez Campus.

Nie, N.H., C.H. Hull, J.G. Jenkins, K. Steinbrenner,

and D-H. Bent. 1975. Statistical Package for the

Social Sciences, 2nd ed. McGraw-Hill Book Company

675. pp.

Peak, M.J. 1983. Killing and mutation by solar uv in

bacteria. Photochem, Photobiol. 37(Suppl.)

PRASA. 1984. Annual report. Available from: PR
Aqueduct and Sewer Authority, San Juan Operationny
Area, San Juan, PR.

Quiftenez-Marquez, P. 1980. Limnology of Lage Loizas
Puerto Rico. US Geological Survey Water Resources
Investigations 79-97. Available from: USGS+ GPO Box
4424, San Juan, PR 00936.

Tuveron, R.W. and ME. March. 1980. Photodynamic and
sunlight inactivation of Escherichia coli strains
Differing in near-vy recombination proficiency,
Photochem. Photobiol. 31(3):267-269.

5, 1977. Topographical map, Sheet 1322 1. stock No.

734%1 3221. Available from? USGS, GPO Box 4424, Sen
van, Ee 00936.

38

---Page Break---

34. velz, C.J. 1970. Applied Stream sanitation. Wiley-
Interscience Publ. Cov, N.Y.

39

---Page Break---

APPENDIX

40.

---Page Break---

Table 4a. Sampling results. NA represents missing valu

Date 3/21/84-Digenal

STP Discharge (m³/hr) 900

Chlorine Applied (kg/hr) 7.56

Parameter

Tine

wig? (ppm {52.0 | 714 | 9.8 | 18.8 | 38.2

a

---Page Break---

Table 4b. Sampling results. NA represents missing value!

Date 3/28/84-Diyrnal

STP Discharge (m³/hr) 792

Chlorine Applied (kg/hr) 9.45

T ~

!

Parameter 1

r T T T

Paro2s 31 #7 5

? 1 1 1 1 1

r r 1 r'

Time 12010 | 1130 | 0945 | 1215 | 1045,

1 ' ' 1'

Fecal Coli. (CFU/mL) 1330 110 128 \$ ae 170

1 1 1 1 1

---Page Break---

Table 4c. Sampling results. NA represents missing values.

Date 4/8/84-Diagnad

STP Discharge (m³/hr) 890

Chlorine Applied (kg/hr) 4.54

T

,

Parameter to

r T

1

riatbobels

epee ds

ng (ppm {93.0 1 73.0 1 110.01 150.01 68.0

---Page Break---

Table 4d. Sampling results. WA represents mi

ing values.

Date 5/2/84-Diurnal

STP Discharge (m³/hr) 792

Chlorine Applied (kg/hr) 5.67

eee

Parameter

T

,

1

,

1

|

1

wm fo | ane |

NHg* (ppm) 1140.01 150.01 280.01 ma na

?4

---Page Break---

Table 4e. Sampling result.

Date \$/9/84-Diagnad

STP Discharge (m4/hr)

Chlorine Applied (ko/he)

Paraneter

Time

Fecal Coli. (CFU/mL)

Log Fecal Coli.

Total coli. (CU/ mt)

Log Total Coli.

Dissolved Oxygen (pa)

Temperature (°C)

Oxygen Saturation (W)

Light Inten. (Candies/ft²)

NO₂⁻/NO₃⁻ (ppm)

Niig* (ppm)

cl⁻ (ppm)

Chlorine (pad)

5. NA represents missing values.

946

Tae

1

i cation

Hi rs rs

' 2) 31 #1 os

I cee Ve

r '' i

10930 | 1008 ; O90 1 1036 | 2045

1 '' i

1115001 8700 | \$60 1 2200 1 300

I ' 1

14.00 i 3.94 1 ».75 13,34 1 2.48

1 1 ? '

1430001 22000; 2300 | 120001 3000

1 ' | 1

14.63 1 4.34 1 5.36 1 4.08 1 3.48

1 1 1 ?

INA TNA fue TNA Ena

1 ''

IWA ENA 1 RA TNA INA

It 1 1 1

INA TNA 1 RR OT NA TNA

1 1 ''

1300 1 2000 1 3000 1 3500 | 1000

'' 1

10.64 1 6.61 | 0.12 | 0.23 1 0.08

? 1 1

148.0 1 150.0} 38.0 1 100.01 68.0

1 1 t 1

Ws 166 150 173 1 69

'' 1 4

Ho.20 1 NA TNR OT NA NA

i 1

---Page Break---

Table 4f. Sampling results. NA represents missing values.

Date 6/20/84-Nockurnal

STP Discharge (m³/hr) 890

Chlorine Applied (kg/hr) 8.98

Parameter

Tine ta0s2

Fecal coli. (cru/m) 14720

log Fecal Coli. ta.aa

Total cous. ccru/my 18500

og otal coli. [3.93

Dissolved Oxygen (ppm) INA

I

Temperature (°C) INA

Oxygen Saturation (9) INA

§02"/1105~ (ppm tent

fag* (ppm) 185.0

c1- (ppm hs

emtorine (ppm) h.s0

46

---Page Break---

Table 4g. Sampling results. NA represents missing values.

Date 6/28/84-Nogturnal

STP Discharge (m/hr) 1003

Chlorine Applied (kg/hr) 3.92

Parameter

47

---Page Break---

Table 4h. Sampling results, NA represents missing values.

Date | 7/2/84-Nockurnal

STP Discharge (mé/hr) 1003

Chlorine Applied (kg/hr) 3.92

Parameter

Terenure 0 fase | asso fate Fane | ae

wry cm hn tan ban fn ba

nay* (ppm 13300 1 2330 1 mn | S650 1 7680

---Page Break---

---Page Break---